

VERIFICATION OF TRANSLATION

I, Naoko Sakai

of 6th Fl., Kantetsu TsukubaScienceCity Bldg. 111, Oroshimachi, Tsuchiura, Ibaraki,
JAPAN

declare as follows:

1. That I am well acquainted with both the English and Japanese languages, and
2. That the attached document prepared by me is, to the best of my knowledge and belief, a true and correct translation of a certified copy of the following application.

Japanese Patent Application No. 2003-374808

Entitled: "METHOD FOR PRODUCING TRANSGENIC DENDRITIC-CELLS"

Filed on November 4, 2003

September 16, 2009
Date

Naoko Sakai
Naoko Sakai

【Name of Document】 APPLICATION FOR PATENT

【Identification Number】 D3-A0307

【Filing Date】 November 4, 2003

【Addressee】 Esq. Commissioner of the Patent Office

【International Patent Classification】 C12N 15/00

【Inventor】

【Address】 15-3-301, Yoshizuka 4-chome, Hakata-ku, Fukuoka-shi, FUKUOKA

【Name】 OKANO, Shinji

【Inventor】

【Address】 5-31-3, Najima, Higashi-ku, Fukuoka-shi, FUKUOKA

【Name】 YONEMITSU, Yoshikazu

【Inventor】

【Address】 1-29-22, Oike, Minami-ku, Fukuoka-shi, FUKUOKA

【Name】 SUEISHI, Katsuo

【Inventor】

【Address】 c/o DNAVEC RESEARCH INC., 25-11, Kannondai 1-chome, Tsukuba-shi,
IBARAKI

【Name】 HASEGAWA, Mamoru

【Applicant】

【Identification Number】 595155107

【Name or Appellation】 DNAVEC RESEARCH INC.

【Attorney】

【Identification Number】 100102978

【Patent Attorney】

【Name or Appellation】 Hatsushi Shimizu

【Nominated Attorney】

【Identification Number】 100108774

【Patent Attorney】

【Name or Appellation】 Kazunori Hashimoto

【Payment】

【Registration Number】 041092

【Amount】 21000 yen

【List of Attached Documents】

【Name of Document】 Claims 1

【Name of Document】 Specification 1

【Name of Document】 Drawings 1

【Name of Document】 Abstract 1

【General Power of Attorney Number】 9716812

[Document Name] Claims

[Claim 1]

A method for producing a gene transferred dendritic cell, which comprises the step of contacting a minus-strand RNA viral vector with a dendritic cell or a precursor cell thereof.

5 [Claim 2]

A method for producing a mature dendritic cell, which comprises the step of contacting a minus-strand RNA viral vector with a dendritic cell or a precursor cell thereof.

[Claim 3]

10 The method of claim 1 or 2, wherein the contacting step involves contacting the minus-strand RNA viral vector with an immature dendritic cell.

[Claim 4]

The method of claim 1 or 2, wherein the contacting step involves contacting the minus-strand RNA viral vector with a CD34⁺ cell.

[Claim 5]

15 The method of claim 3 or 4, further comprising the step of culturing the cell in the presence of GM-CSF and IL-4 before or after the contact step.

[Claim 6]

20 The method of claim 1 or 2, wherein the minus-strand RNA viral vector is a paramyxovirus vector.

[Claim 7]

The method of claim 1 or 2, wherein the paramyxovirus vector is a Sendai virus vector.

[Claim 8]

The method of claim 1 or 2, wherein the cell is a human cell.

25 [Claim 9]

A vector-comprising cell produced by the method of any one of claims 1 to 8.

[Claim 10]

The cell of claim 9, which is a mature dendritic cell.

[Document Name] Specification

[Title of the Invention] METHOD FOR PRODUCING TRANSGENIC DENDRITIC-CELLS

[Technical Field]

[0001]

5 The present invention relates to a method for introducing genes into dendritic cells. The methods of the present invention can be used in the production of vaccines against cancers, infectious diseases, and the like.

[Background Art]

[0002]

10 The dendritic cell (DC) is one of the antigen-presenting cells (APCs) present in peripheral blood, skin, lymphatic organs, thymus, and such, and is widely distributed in lymphatic and non-lymphatic tissues (see Steinman, R. M., Ann. Rev. Immunol. (1991) 9: 271; Banchereau, J.B. and Steinman R.M., Nature (1998) 392: 245). The dendritic cell has strong antigen-presenting ability and expresses antigen peptides on class I and II on the dendritic cell
15 surface, which activate CD4 and CD8 T cells, respectively. Through this activation, the cell induces an *in vivo* immune response against specific antigens (e.g., antigens of pathogenic microorganisms, tumor-related antigens, transplantation antigens, *etc.*).

[0003]

 Gene alterations in dendritic cells produce a variety of clinically beneficial effects. For
20 example, the production and use of a mature dendritic cell expressing a molecule (CD80, CD86, and the like) that induces a costimulatory signal required for T cell response enables strong activation of adaptive immunity against an antigen (for example, vaccines against viruses and tumors). Alternatively, immune tolerance to an antigen (for example, acquired insensitivity to transplantation antigens and antigens causing autoimmune diseases) can be induced by using
25 dendritic cells devoid of a costimulatory signal-inducing molecule or a dendritic cell expressing a molecule that gives suppressive stimulus. Reported examples of such applications include: immunostimulation by introducing the interleukin (IL)-12 gene into dendritic cells (Gene Therapy 7: 2113-2121 (2000)), and removal of antigen specific T cells by introducing the Fas ligand gene into dendritic cells (J Immunol. 200: 164; 161-167). With respect to tumor
30 immunotherapy, immunity against tumor can be induced through gene transfer of a tumor antigen into dendritic cells.

[Non-Patent Document 1] Steinman, R. M., 1991, Ann. Rev. Immunol. 9: 271

[Non-Patent Document 2] Banchereau, J.B. and R.M. Steinman, 1998, Nature 392: 245

[Non-Patent Document 3] Akiyama, Y. et al., 2000, Gene Therapy, 7: 2113-2121

35 [Non-Patent Document 4] Min, W.P., 2000, J. Immunol., 164: 161-167

[Disclosure of the Invention]

[Problems to Be Solved by the Invention]

[0004]

The present invention provides a method for introducing genes into dendritic cells.

- 5 The present invention also provides a method for producing gene transferred dendritic cells. Furthermore, the present invention provides dendritic cells that are introduced with genes by the methods of the present invention. In addition, the present invention provides a viral vector for gene transfer into dendritic cells. The present invention also provides a method for activating dendritic cells.

10 [Means for Solving the Problems]

[0005]

- The present inventors discovered that minus-strand RNA viral vector is an excellent vector for gene transfer into dendritic cells. Sufficient gene transfer efficiency can be achieved through only a short contact time between the minus-strand RNA viral vector and dendritic cells, and the expression was detectable for a long period of time. Furthermore, dendritic cells can be activated through mere infection with the minus-strand RNA viral vector.

- Although gene transfer into dendritic cells is expected to be applicable to various types of immunotherapy, previous techniques for gene transfer into dendritic cells were complicated and/or their introduction efficiency was unsatisfactory. The present invention has demonstrated that foreign genes can be introduced into dendritic cells by a very simple procedure using a minus-strand RNA viral vector. Efficient gene delivery into dendritic cells can be achieved by the method of the present invention, and thus the method is expected to be applicable to gene alteration of dendritic cells for immunotherapy.

[0006]

- 25 Specifically, the present invention relates to a method for introducing genes into dendritic cells, and the like, and, more specifically, relates to:
- [1] a method for producing a gene transferred dendritic cell, which comprises the step of contacting a minus-strand RNA viral vector with a dendritic cell or a precursor cell thereof;
 - [2] a method for producing a mature dendritic cell, which comprises the step of contacting a minus-strand RNA viral vector with a dendritic cell or a precursor cell thereof;
 - 30 [3] the method of [1] or [2], wherein the contacting step involves contacting the minus-strand RNA viral vector with an immature dendritic cell;
 - [4] the method of [1] or [2], wherein the contacting step involves contacting the minus-strand RNA viral vector with a CD34⁺ cell;
 - 35 [5] the method of [3] or [4], further comprising the step of culturing the cell in the presence of GM-CSF and IL-4 before or after the contact step;

[6] the method of [1] or [2], wherein the minus-strand RNA viral vector is a paramyxovirus vector;

[7] the method of [1] or [2], wherein the paramyxovirus vector is a Sendai virus vector;

[8] the method of [1] or [2], wherein the cell is a human cell;

5 [9] a vector-comprising cell produced by the method of any one of [1] to [8]; and

[10] the cell of [9], which is a mature dendritic cell.

[Effects of the Invention]

[0007]

Dendritic cells have high ability for immune induction. Thus, dendritic cell (DC)
10 vaccines useful for immunotherapy against cancers, infectious diseases, and the like can be produced by introducing a desired antigen gene or an immune activating gene into dendritic cells through the present method. For example, to present tumor antigens on dendritic cells for tumor immunotherapy, a method of mixing dendritic cells with the lysate of tumor cells, a peptide pulse method, a method of introducing tumor antigen genes into dendritic cells, and the
15 like may be used. Among them, the method of introducing tumor antigen genes into dendritic cells can be expected to prolong the duration of *in vivo* tumor antigen presentation as compared with the tumor lysate and peptide pulse methods, and also has the advantage of being free from HLA restriction (in the case of peptide: a certain peptide derived from an antigen is used; however, due to the requirement for the binding with HLA, the region of the antigen used for the
20 peptide varies depending on the type of HLA).

[0008]

The liposome method, electroporation, and the like are available to introduce plasmids as vectors for introducing genes into dendritic cells. However, these methods are considered to be impractical due to their low introduction efficiency (Cancer Gene Ther 4: 17-25s (1997)).
25 Practical vectors include the following three kinds of vectors. (i) adenoviral vectors (J. Immunotherapy 25: 445-454 (2002); Gene Therapy 7: 249-254 (2000)), (ii) retroviral vectors (J. Leuko. Biol., 263-267 (1999); Br. J. Haematol. 108: 817-824 (2000)), and (iii) lentiviral vectors (J. Gene Med. 3: 311-320 (2001); J. Immunol. Meth. 153-165 (2002); Mol. Ther. 283-290 (2002); Cancer Gene Therapy 9: 715-724 (2002)).

30 [0009]

Of the above, the retroviral vectors of (ii) can only be introduced into cells of the growth phase. In addition, these vectors require cytotoxic reagents, such as polybrene, that assist their introduction and their introduction is substantially time consuming. Therefore, they are disadvantageous in that they tend to lower the cell viability. Furthermore, the introduction
35 efficiency of these vectors into dendritic cells induced from peripheral blood is much worse. Thus, in general, the differentiation of the cells to dendritic cells is induced after the introduction

of the vectors into CD34-positive cells. Since this method requires bone marrow cells, cord blood, or peripheral blood immobilized with G-CSF, it is very invasive to patients. In addition, the introduction of retroviral vectors into dendritic cells fails to elevate the activation state of dendritic cells. Furthermore, retroviral vectors integrate viral nucleic acids into the genome of the cells, which raises concerns about damage to the genome. Moreover, the introduction efficiency is generally low, and thus requires selection of the vector-transferred cells via sorting or the like.

[0010]

Vector introduction using the lentiviral vectors of (iii), like retroviral vectors, is very time consuming. This, in turn, can lower cell viability. The lentivirus is also known to transfer genes into resting cells. However, in general, due to postentry restriction, the efficiency achieved by the introduction into dendritic cells that are differentiated to some extent and have lost their proliferating activity is very low (several %) (J. Virol. 75; 5448-5456). Furthermore, the introduction efficiency into dendritic cells induced from peripheral blood is much lower. Therefore, the lentiviral vectors can only be used through a method wherein dendritic cells are obtained *in vitro* after the introduction of vectors into CD34 positive stem cells (this method is quite invasive to patients since it requires bone marrow cells, cord blood, or peripheral blood immobilized with G-CSF). Recent vector modification techniques attempt to overcome this problem, and, today, in the case of SIV, gene transfer into peripheral blood-derived monocytes and differentiated dendritic cells can be achieved by retaining vpx (which promotes nuclear localization of proviral DNA) in the helper construct or by inserting a DNA-flap sequence (which also promotes nuclear localization of proviral DNA) with HIV (Mol. Ther. 283-290 (2002)). However, it is still necessary to infect the virus at an early stage of the differentiation of monocytes to dendritic cells and therefore, the differentiation of the dendritic cells may be disturbed by the inserted gene. In addition, the introduction efficiency varies from donor to donor. Furthermore, the introduction of vector cannot elevate the activation state of the dendritic cells. Similarly to the retroviral vector described above, use of the lentiviral vectors involve risks of genome damage and development of malignant tumors, which is a barrier to practical use of the vector at present. Moreover, due to the low introduction efficiency, selection is generally required or higher titers of the vectors are required to improve the introduction efficiency, which hampers the clinical application of the vectors. In addition, the vector requires auxiliary manipulations (such as centrifugation) or assisting reagents (such as polybrene), as well as longer reaction time due to the complicated introduction procedures. The cost needed for vector purification is an additional problem.

[0011]

Meanwhile, due to the introduction efficiency (about 80%) and its ability to directly

transfer genes into differentiated cells, the adenovirus of (i) is the currently preferred vector for gene transfer into dendritic cells. However, the introduction efficiency is time-dependent and it requires about 72 hours to gain maximal introduction efficiency (J. Immunotherapy 25: 445-454 (2002)). Furthermore, this vector requires 10 to 100 times higher titer as compared with the minus-strand RNA virus. A more critical problem is that this vector, at MOIs allowing high gene transfer efficiency, has an immunosuppressive effect, which attenuates the mixed lymphocyte reaction (MLR) of allo T cells (in particular, at high DC:T ratios) (Gene Therapy 2000; 7; 249-254). In addition, due to dilution of episome, it is sometimes difficult to perform the step for differentiating dendritic cells from stem cells, such as CD34 positive cells, after gene transfer.

[0012]

Requirements to achieve gene introduction into dendritic cells include high introduction efficiency, technical stability for gene transfer, convenience, clinical safety, and maintenance of the T cell activating ability of the dendritic cells. However, none of the presently available vectors satisfy all these requirements. In contrast, when the minus-strand RNA viral vector was used, a very short contact period resulted in a gene introduction efficiency of nearly 100% and the suppression of allo T cell response was relatively mild to retain the T cells stimulating ability of the cell. Thus, the minus-strand RNA viral vector appears to meet all requirements described above, considering that the vector has the ability to express proteins in the cytoplasm, is safe, and requires neither nuclear transfer nor gene insertion to the genome. The gene transfer of the vectors of (i) to (iii) described above do not alter the activation state of dendritic cells. In contrast, the gene transfer of the minus-strand RNA viral vector induces dendritic cell activation, and thus, when used for immunostimulation (e.g., tumor immunity, *etc.*), the post-transfer step for the activation with cytokines and the like can be omitted, which is expected to contribute to maintenance of cell viability, reduction in cost, and further reduction in the time required for *ex vivo* manipulation. It was also herein confirmed that activated T cells, in particular, tumor specific cytotoxic T cells and the like, required for T cell transfer therapy could be efficiently and easily induced *ex vivo* in a short period by using dendritic cells gene transferred with the minus-strand RNA viral vector. In addition, comparable to the lentiviral vectors, when dendritic cells were differentiated after the gene transfer of the vector into stem cells, the introduction efficiency reached nearly about 70%. These characteristics expand the scope of clinical applications of the minus-strand RNA viral vector.

[Best Mode for Carrying Out the Invention]

[0013]

The present invention provides a method for producing dendritic cells introduced with genes using a minus-strand RNA viral vector. This method includes the step of contacting a

minus-strand RNA viral vector carrying a gene desired to be introduced with a dendritic cell or a precursor cell thereof. The present inventors discovered that the minus-strand RNA viral vector was capable of introducing a gene into dendritic cells with very high efficiency. In addition, the introduction of the vector activated the dendritic cell leading to differentiation into mature
5 dendritic cell without cytokine stimulation. The resulting mature dendritic cell retained the ability to activate T cells. The present method is useful for the presentation of desired antigens by dendritic cells, or to express desired cytokines or other physiologically active factors in the dendritic cells. Dendritic cells that are genetically modified by the present method have high ability to activate the immune system, and can be suitably used to prevent or treat infectious
10 diseases, cancers, and other desired diseases on which immune induction is expected to cause beneficial effects. The contact between the vector and dendritic cells can be achieved *in vivo* or *in vitro*, for example, in a desired physiological aqueous solution, such as culture solution, physiological saline, blood, blood plasma, serum, and body fluid.

[0014]

15 The introduction efficiency of the minus-strand RNA viral vector is significantly higher in non-activated (immature) dendritic cells than in mature dendritic cells. Thus, it is preferred to contact the minus-strand RNA viral vector with immature dendritic cells or to mix the vector with a cell fraction containing immature dendritic cells. Such methods are also included in the method of the present invention for introducing genes into dendritic cells. Dendritic cells are
20 activated by the contact with bacteria, lipopolysaccharide (LPS), double-stranded RNA, or the like. When dendritic cells to be introduced with genes are separately activated by such a method, to prevent the reduction in the efficiency of vector introduction, the activation treatment is preferably carried out not before the vector introduction but after gene transfer using the minus-strand RNA viral vector (or simultaneous with the contact of the minus-strand RNA viral
25 vector to the dendritic cells).

[0015]

The achievement of highly efficient gene delivery through simple techniques is an important superiority of the minus-strand RNA viral vector-mediated gene delivery to the dendritic cells. The gene delivery into dendritic cells mediated by retroviral vectors and the
30 like has low efficiency, and sometimes requires a toxic agent, such as polybrene, to stimulate gene transfer. On the other hand, superior gene delivery can be achieved with the minus-strand RNA viral vector, without any requirement for a special agent, by simply adding to a solution containing dendritic cells. Furthermore, the minus-strand RNA viral vector-mediated gene delivery into dendritic cells can achieve the highest efficiency within a very short exposure time
35 (for 30 minutes or less). Considering clinical situations, these characteristics simplify *ex vivo* and *in vivo* genetic alterations of dendritic cells, and can minimize manipulation-dependent

adverse effects, such as loss of cell viability.

[0016]

For the contact of the vector with dendritic cells, MOI (multiplicity of infection: the number of infecting viruses per cell) is preferably within the range of 1 to 500, more preferably within the range of 2 to 300, even more preferably within the range of 3 to 200, still more preferably within the range of 5 to 100, and yet more preferably within the range of 7 to 70. The contact between the vector and dendritic cells requires only a short time, which may be, for example, 1 minute or longer, preferably 3 minutes or longer, 5 minutes or longer, 10 minutes or longer, or 20 minutes or longer, for example, within the range of about 1 to 60 minutes, more specifically within the range of about 5 to 30 minutes. Of course, the contact time may be longer, for example, for several days or longer. The contact can be achieved *in vivo* or *ex vivo*. For example, the present method is suitably used in *ex vivo* gene transfer where dendritic cells or precursor cells thereof removed from the body are contacted *ex vivo* with the minus-strand RNA viral vector, and returned into the body after vector introduction.

[0017]

One of the characteristics of the present method is the sustained long-term expression of introduced gene in dendritic cells after the gene transfer. According to the present method, the expression of the introduced gene is detected in the dendritic cells for 2 days or more, for example, for 3 days or more, 5 days or more, 10 days or more, 14 days or more, 30 days or more, 50 days or more, and even 60 days or more after the infection of the vector into the cells.

[0018]

The present invention also provides a method for selectively introducing a gene into immature dendritic cells, which includes the step of allowing the minus-strand RNA viral vector carrying the gene to coexist with a cell population composed of mature and immature dendritic cells. The phrase "selectively ... into immature dendritic cells" means that the gene is introduced into immature dendritic cells at a significantly high rate as compared with mature dendritic cells. Specifically, the ratio of the vector-introduced immature dendritic cells to the total immature dendritic cells is significantly higher than the ratio of the vector-introduced mature dendritic cells to the total mature dendritic cells. For example, the present invention provides a method which includes the step of adding a minus-strand RNA viral vector carrying a desired gene to be introduced to a cell population composed of mature and immature dendritic cells. Since the minus-strand RNA viral vector preferentially transfers genes into immature dendritic cells as compared to mature dendritic cells, a gene can be selectively introduced into immature dendritic cells through this method. Such methods are also included in the method of the present invention for introducing genes into dendritic cells.

[0019]

A dendritic cell (DC) is a cell which takes a dendritic morphology in the mature state and has the ability to activate T cells by presenting an antigen. The dendritic cells include a group of bone marrow-derived cells with dendritic morphology distributed in various organs and tissues in the body, and a group of cells resulting from *in vitro* differentiation of bone marrow- or blood-derived stem cells using cytokines or the like, that are equivalent to the cells with dendritic morphology distributed in various organs and tissues in the body. Specifically, the dendritic cells include, for example, lymphocytic dendritic cells (including cells which induce Th2 or immune tolerance), bone marrow dendritic cells (generally used dendritic cells, including immature and mature dendritic cells), Langerhans cells (dendritic cells important as antigen-presenting cells in the skin), interdigitating cells (distributed in the lymph nodes and spleen T cell region, and believed to function in antigen presentation to T cells), and follicular dendritic cells (important as antigen-presenting cells for B cells; the cells present antigens to B cells by presenting antigen-antibody complexes or antigen-complement complexes on the surface via the antibody receptor or the complement receptor). Preferably, the dendritic cells highly express MHC class I and class II, and more preferably express CD11c.

[0020]

A dendritic cell may also be a cell with dendritic morphology and that is positive for two or more surface markers selected from the group consisting of CD11c, HLA-class II (HLA-DR, -DP, or -DQ), CD40, and CD1a. The dendritic cell of the present invention is more preferably an HLA-class II⁺ and CD11c⁺ cell, even more preferably, a CD1a⁺, HLA-class II⁺, and CD11c⁺ cell that is devoid of the expression of T cell marker (CD3), B cell markers (CD19, CD20), NK cell marker (CD56), neutrophil marker (CD15), and monocyte marker (CD14). The proportion of CD14⁺ cells in a dendritic cell population to be used for the vector introduction is, for example, 10% or less, preferably 5% or less, and more preferably 1% or less.

[0021]

In addition, the dendritic cells of the present invention include both mature and immature dendritic cells. The immature dendritic cells refer to dendritic cells having low antigen-presenting ability for T cell activation and the like, preferably having phagocytic ability for antigens, and more preferably showing low or negative expression of receptors that induce the costimulation for T cell activation. On the other hand, the mature dendritic cells refer to dendritic cells that have strong antigen-presenting ability for T cell activation or the like, preferably have weak or no phagocytic ability for antigen, and more preferably show high expression of receptors that induce the costimulation for T cell activation. The activation of dendritic cells refers to the transition from immature to mature dendritic cell; and the activated dendritic cells encompass mature dendritic cells and dendritic cells in the process of the transition, wherein the expression of CD80 and CD86 that induce costimulatory signals are

elevated upon the activating stimuli. In CD11c positive dendritic cells, CD83 positivity serves as an indicator of mature dendritic cells.

[0022]

For example, mature dendritic cells may preferably be cells which highly express CD40, CD80, CD86, and HLA-class II. More preferably, mature dendritic cells express CD83. An immature dendritic cell can be distinguished from a mature dendritic cell, for example, using markers selected from the group consisting of CD80, CD83, and CD86. The immature dendritic cell is weakly positive or negative (dim/negative) for these markers, while the mature dendritic cell is positive.

[0023]

Furthermore, immature dendritic cells generally have a high phagocytic ability. When dendritic cells are added with LPS (1 μ g/ml) and cultured for two days, they become activated and their phagocytic ability is reduced. The phagocytic ability can be detected by measuring the amount of small molecules taken up into dendritic cells or the proportion of uptaking cells. The phagocytic ability is preferably determined by the amount of small molecules taken up into dendritic cells. For example, using colored beads with a size of about 1 μ m, the uptake of beads into dendritic cells can be measured. Quantitation is performed by subtracting the positive background at 4°C. Among obtained dendritic cell groups, dendritic cell group with high phagocytic ability indicates an ability wherein the amount of small molecules taken up into dendritic cells is 4 times or more, more preferably 5 times or more, and even more preferably 6 times or more than that taken up into dendritic cells stimulated with LPS for two days as described above. Alternatively, the proportion of cells taking up small molecules is twice or more, and more preferably 3 times or more.

[0024]

The respective markers described above and methods for measuring their expression are also well known to those skilled in the art. For example, CD11c is an adhesion glycoprotein of about 150 kD (p150, integrin alpha chain). CD11c binds to CD18 to form a CD11c/CD18 complex, which is capable of binding to fibrinogen and has been reported to function as a receptor for iC3b and ICAM-1. In addition, it has been reported that CD11c/CD18 can function as an adhesion molecule that binds to receptors on stimulated epithelia (Knapp, W. *et al.*, eds., 1989, Leucocyte Typing IV: White Cell Differentiation Antigens, Oxford University Press, New York; Barclay, N.A. *et al.*, eds., 1993, The Leucocyte Antigen Facts Book, CD11 Section, Academic Press Inc., San Diego, California, p. 124; Stacker, S.A. and T.A. Springer, 1991, J. Immunol. 146: 648).

[0025]

CD1a is a polypeptide of about 49 kD, which binds to beta2 microglobulin. CD1a is

structurally similar to an MHC class I antigen and is assumed to function in antigen presentation (Knapp, W. *et al.*, eds., 1989, Leucocyte Typing IV: White Cell Differentiation Antigens, Oxford University Press, New York; Schlossman, S. *et al.*, eds., 1995, Leucocyte Typing V: White Cell Differentiation Antigens. Oxford University Press, New York; Hanau, D. *et al.*, 1990, J.

- 5 Investigative Dermatol. 95: 503; Calabi, F. and A. Bradbury., 1991., Tissue Antigens 37: 1).
[0026]

CD14 is a glycosylphosphatidylinositol (GPI)-anchored single-chain glycoprotein of 53 to 55 kD expressed in dendritic reticulum cells and some types of Langerhans cells. CD14 was identified as a surface receptor having high affinity to a complex of LPS and serum LPS-binding protein (LPB) (McMichael, A.J. *et al.*, eds., 1987, Leucocyte Typing III: White Cell Differentiation Antigens, Oxford University Press, New York; Knapp, W. *et al.*, eds., 1989, Leucocyte Typing IV: White Cell Differentiation Antigens, Oxford University Press, New York; Schlossman, S. *et al.*, eds., 1995, Leucocyte Typing V: White Cell Differentiation Antigens. Oxford University Press, New York; Wright, S.D. *et al.*, 1990, Science 249: 1434).

- 15 [0027]

CD40 is a type I integral membrane protein of 45 to 48 kD (type I integral membrane glycoprotein). Anti-CD40 antibody is frequently used as a cell marker (Schlossman, S. *et al.*, eds., 1995, Leucocyte Typing V: White Cell Differentiation Antigens. Oxford University Press, New York; Galy, A.H.M.; and H. Spits, 1992, J. Immunol. 149: 775; Clark, E.A. and J.A.

- 20 Ledbetter, 1986, Proc. Natl. Acad. Sci. 83: 4494; Itoh, H. *et al.*, 1991, Cell 66: 233; Barclay, N.A. *et al.*, 1993, The Leucocyte Antigen Facts Book., Academic Press).

CD80 is a transmembrane glycoprotein of about 60 kD, and is a member of the Ig supergene family. CD80 is a ligand for CD28 and CD152 (CTLA-4) expressed in T cells (Schlossman, S. *et al.*, eds., 1995, Leucocyte Typing V: White Cell Differentiation Antigens.

- 25 Oxford University Press, New York; Schwartz, R.H., 1992, Cell 71: 1065; Azuma, M. *et al.*, 1993, J. Exp. Med. 177: 845; Koulova, L. *et al.*, 1991, J. Exp. Med. 173: 759; Freeman, G.J. *et al.*, 1998, J. Immunol. 161: 2708; Behrens, L. *et al.*, 1998, J. Immunol., 161(11):5943; Guesdon, J.-L. *et al.*, 1979, J. Histochem. Cytochem. 27: 1131-1139).

[0028]

30 CD83 is a transmembrane protein of about 45 kD, and is a member of the Ig superfamily. CD83 has a short extracellular domain of V-type Ig and a C-terminal cytoplasmic tail. CD83 is mainly expressed in follicular dendritic cells, circulating dendritic cells, interdigitating dendritic cells in lymphatic tissues, *in vitro*-produced dendritic cells, and dendritic cells of the thymus (Zhou, L.-J., and T.F. Tedder, 1995, J. Immunol. 154: 3821; Zhou, L.-J. *et al.*, 1992, J. Immunol. 149: 735; Summers, K.L. *et al.*, 1995, Clin Exp. Immunol. 100: 81; Weissman, D. *et al.*, 1995, Proc. Natl. Acad. Sci USA. 92: 826; Hart, D.N.J., 1997, Blood 90: 3245).

- 35

CD86 (B70/B7-2) is a cell surface protein of about 75 kD, which is a second ligand for CD28 and CTLA-4 and plays an important role in costimulation of T cells in early immune response (Azuma M. *et al.*, 1993, *Nature* 366: 76; Nozawa Y. *et al.*, 1993, *J. Pathology* 169: 309; Engle, P. *et al.* 1994., *Blood* 84: 1402; Engel, P. *et al.*, CD86 Workshop Report. In: Leukocyte
 5 Typing V. Schlossman, S.F. *et al.* eds., 1994, Oxford University Press; Yang, X.F. *et al.*, 1994, Upregulation of CD86 antigen on TPA stimulated U937 cells, 1994, (abstract). American Society of Hematology, Nashville, TN; Guesdon, J.-L. *et al.*, 1979, *J. Histochem. Cytochem.* 27: 1131-1139).

[0029]

10 CCR7 is also called BLR-2, EBI-1, and CMKBR7, which is a seven-transmembrane G protein-coupled receptor, and is a receptor of the CC chemokines, MIP-3beta/Exodus 3/ELC/CCL19 and 6Ckine/Exodus 2/SLC/TCA4/CCL21 (Sallusto, F. *et al.*, 1999, *Nature* 401: 708-12; Lipp, M. *et al.*, 2000, *Curr. Top. Microbiol. Immunol.* 251: 173-9; Birkenbach, M. *et al.*, 1993, *J. Virol.* 67: 2209-20; Schweickart, V. L. *et al.*, 1994, *Genomics* 23: 643-50; Burgstahler, R.
 15 *et al.*, 1995, *Biochem. Biophys. Res. Commun.* 215: 737-43; Yoshida, R. *et al.*, 1997, *J. Biol. Chem.* 272:13803-9; Yoshida, R. *et al.*, 1998, *J. Biol. Chem.* 273: 7118-22; Yoshida, R. *et al.*, 1998, *Int. Immunol.* 10: 901-10; Kim, C. H. *et al.*, 1998, *J. Immunol.* 161: 2580-5; Yanagihara, S. *et al.*, 1998, *J. Immunol.* 161: 3096-102).

[0030]

20 DR, DP, and DQ exist as HLA-class II antigens, and can be collectively detected using antibodies that bind to all three antigens (Pawelec, G. *et al.*, 1985, *Human Immunology* 12: 165; Ziegler, A. *et al.*, 1986, *Immunobiol.* 171: 77). HLA-DR is one of the human MHC class II antigens, which is a transmembrane glycoprotein consisting of an alpha chain (36 kDa) and a beta subunit (27 kDa). In epidermal Langerhans cells, the protein is co-expressed with CD1a
 25 antigen. CD1a plays a principal role in cell interaction for antigen presentation (Barclay, N.A. *et al.*, 1993, *The Leucocyte Antigen Facts Book*. p. 376. Academic Press).

[0031]

Dendritic cells of nonhuman mammals can also be specified using the products of homologous genes of the above-described marker genes as indicators. Antibodies to such
 30 markers are commercially available, for example, from BD Biosciences (BD PharMingen), and detailed information is available at the websites of the company or its distributors.

[0032]

For dendritic cell markers, also see the references by Kiertscher *et al.* and Oehler *et al.*. (Kiertscher SM, Roth MD, Human CD14⁺ leukocytes acquire the phenotype and function of
 35 antigen-presenting dendritic cells when cultured in GM-CSF and IL-4, *J. Leukoc. Biol.*, 1996, 59 (2): 208-18; Oehler, L. *et al.*, Neutrophil granulocyte-committed cells can be driven to acquire

dendritic cell characteristics., J. Exp. Med., 1998, 187 (7): 1019-28). For further details regarding flow cytometry, see the references by Okano *et al.* and Stites *et al.* (Okano, S. *et al.*, Recombinant Sendai virus vectors for activated T lymphocytes. Gene Ther., 2003, 10(16): 1381-91; Stites, D. *et al.*, Flow cytometric analysis of lymphocyte phenotypes in AIDS using monoclonal antibodies and simultaneous dual immunofluorescence., Clin. Immunol. Immunopathol., 1986, 38: 161-177). The expression of each of the markers may be determined, for example, using as a threshold the fluorescence intensity that makes a positive rate of 1% or less when stained with an isotype control antibody, wherein the fluorescence equal to or above the threshold is deemed positive, and the fluorescence below deemed negative.

10 [0033]

Dendritic cells or precursor cells thereof can be prepared according to or based on known methods. For example, the cells can be isolated from blood (for example, peripheral or cord blood), bone marrow, lymph nodes, other lymphatic organs, spleen, and skin. Dendritic cells to be used in the context of the present invention are preferably obtained from blood or bone marrow. Alternatively, dendritic cells to be used in the present invention may be skin Langerhans cells, veiled cells of afferent lymphatics, follicular dendritic cells, spleen dendritic cells, and interdigitating cells of lymphatic organs. The dendritic cells used in the present invention include dendritic cells selected from the group consisting of CD34⁺-derived dendritic cells, bone marrow-derived dendritic cells, monocyte-derived dendritic cells, splenic cell-derived dendritic cells, skin-derived dendritic cells, follicular dendritic cells, and germinal center dendritic cells. CD34⁺-derived dendritic cells can be differentiated from hematopoietic stem cells, hematopoietic progenitor cells, or the like, obtained from cord blood, bone marrow, or the like, using granulocyte colony stimulating factor (G-CSF), granulocyte macrophage colony stimulating factor (GM-CSF), tumor necrosis factor (TNF)-alpha, IL-4, IL-13, stem cell factor (SCF), Flt-3 ligand, c-kit ligand, combinations thereof, or the like. For example, peripheral blood monocytes can be differentiated into immature dendritic cells using GM-CSF and IL-4, and then differentiated into mature dendritic cells by stimulating with TNF-alpha.

[0034]

When dendritic cells are selected (or enriched) from a composition including dendritic cells and other cells, it is preferable to perform so-called negative selection which removes cells other than the dendritic cells. Through the negative selection process, precursors of DC-granulocytes (J. Exp. Med., 1998, 187: 1019-1028; Blood, 1996, 87: 4520-4530) remain and thus, it is considered that not only DCs differentiated from adhesive CD14⁺ cells but also DCs differentiated from precursors can be recovered together. This is expected to reduce the cytotoxicity resulting from vector introduction.

[0035]

For example, by removing T cells, NK cells, B cells, and the like, using antibodies specific thereto, dendritic cells can be enriched. Specifically, for example, it is preferable to obtain cells with low or negative expression of a surface marker selected from CD2, CD3, CD8, CD19, CD56, and CD66b, or any combinations thereof. More preferred are cells in which the expressions of CD2, CD3, CD8, CD19, CD56, and CD66b are all low or negative. Therefore, it is preferable to remove cells expressing these markers using antibodies against the markers (Hsu *et al.*, Nature Med. 2: 52 (1996)). The negative selection can be performed using polyvalent antibodies as shown in the Examples. Alternatively, a similar selection can also be performed using beads or the like for magnetic cell separation (MACS). The use of beads is preferred for large scale cell preparation, such as collection of mononuclear cells through blood cell separation or the like. For example, dendritic cells prepared by negative selection from monocytes that were enriched from a cell solution obtained from the body can be preferably used in the context of the present invention.

[0036]

When dendritic cells differentiated from peripheral blood monocytes obtained from adhesive cells are selected before introduction of the minus-strand RNA virus, the efficiency of vector introduction is sometimes reduced. To prevent the reduction of the proportion of immature dendritic cells, before the contact with the minus-strand RNA viral vector, cell culture is preferably carried out without the step of selecting cells adhering to a solid support (for example, culture container such as culture dish or bottle); however, the dendritic cells used in the context of the present invention are not limited thereto. Specifically, the present invention provides a method which excludes the step of selecting cells adhered on the solid support within 24 hours before contact of dendritic cells with the minus-strand RNA viral vector. More preferably, the method excludes the step of selecting cells adhered to the solid support within 2, 3, 5, or 7 days before the contact of dendritic cells with the minus-strand RNA viral vector.

[0037]

The method preferably excludes the step of selecting CD14⁺ cells before the contact with the minus-strand RNA viral vector, but it is not limited thereto. Specifically, the present invention provides a method which excludes the step of selecting CD14⁺ cells within 24 hours before the contact of dendritic cells with the minus-strand RNA viral vector. More preferably, the method excludes the step of selecting CD14⁺ cells within 2, 3, 5, or 7 days before the contact of dendritic cells with the minus-strand RNA viral vector.

[0038]

Specific methods for isolating dendritic cells are described in, for example, Cameron *et al.*, Science 257: 383 (1992); Langhoff *et al.*, Proc. Natl. Acad. Sci. USA 88: 7998 (1991); Chehimi *et al.*, J. Gen. Virol. 74:1 277 (1993); Cameron *et al.*, Clin. Exp. Immunol. 88: 226

(1992); Thomas *et al.*, 1993, J. Immunol. 150: 821 (1993); and Karhumaki *et al.*, Clin. Exp. Immunol. 91: 482 (1993). The isolation of dendritic cells by flow cytometry is described in, for example, Thomas *et al.*, J. Immunol. 153: 4016 (1994); Ferbas *et al.*, J. Immunol. 152: 4649 (1994); and O'Doherty *et al.*, Immunology 82: 487 (1994). In addition, magnetic cell separation is described in, for example, Miltenyi *et al.*, Cytometry 11: 231-238 (1990).

[0039]

Furthermore, for example, human dendritic cells may be isolated and grown using the methods described in Macatonia *et al.*, Immunol. 74: 399-406 (1991); O'Doherty *et al.*, J. Exp. Med. 178: 1067-1078 (1993); Markowicz *et al.*, J. Clin. Invest. 85: 955-961 (1990); Romani *et al.*, J. Exp. Med. 180: 83-93 (1994); Sallusto *et al.*, J. Exp. Med. 179: 1109-1118 (1994); Bernhard *et al.*, J. Exp. Med. 55: 1099-1104 (1995); and the like. Moreover, dendritic cells can be formed from CD34⁺ cells obtained from bone marrow, cord blood, peripheral blood, or the like and from peripheral blood-derived mononuclear cells by the method described in Van Tendeloo *et al.*, Gene Ther. 5: 700-707 (1998).

[0040]

In the present invention, it is preferable to mix a minus-strand RNA viral vector with a cell fraction containing a high density of dendritic cells or precursor cells thereof (for example, CD11c⁺ cells or CD34⁺ cells). The precursor cells refer to cells that can differentiate into dendritic cells in the presence of appropriate cytokines. The precursor cells are preferably differentiated into dendritic cells within 4 weeks, more preferably within 20 days, even more preferably within 18 days, and still more preferably within 16 days. Such cells include CD34⁺ stem cells. The differentiation into dendritic cells may be achieved, for example, by culturing the cells in the presence of SCF (50 ng/ml), GM-CSF (500 U/ml), and TNF-alpha (50 ng/ml) for about 3 days, followed by culturing in the presence of SCF (50 ng/ml), GM-CSF (500 U/ml), IL-4 (250 U/ml), and TNF-alpha (50 ng/ml). A cell fraction refers to a group of cells obtained through cell separation (or cell fractionation). The cell fraction may be a composition including both cells and pharmaceutically acceptable carriers. Exemplary carriers include desired solutions that can be used to suspend viable cells, such as physiological saline, phosphate buffered saline (PBS), culture medium, and serum. According to the present method, the cell fraction to be contacted with the vector includes dendritic cells and/or precursors thereof at a proportion of, for example, 30% or more, preferably 40% or more, preferably 50% or more, preferably 60% or more, and preferably 70% or more to the total viable cells.

[0041]

Furthermore, dendritic cells to be contacted with a minus-strand RNA viral vector preferably include immature dendritic cells. In the cell fraction that includes dendritic cells to be mixed with the vector, the proportion of immature dendritic cells to the total viable cells is,

for example, 10% or more, preferably 20% or more, more preferably 30% or more, more preferably 40% or more, more preferably 50% or more, more preferably 60% or more, and more preferably 70% or more.

[0042]

5 The method of the present invention using a minus-strand RNA virus has various advantages. For example, with the minus-strand RNA virus, activated dendritic cells can be obtained by the vector infection alone, and subsequent steps for obtaining mature dendritic cells can be omitted. Since dendritic cells must be activated for their use in immunostimulation, it is advantageous that the activation can be achieved by only the vector infection. Furthermore, 10 utilizing this property, activated T cells, in particular cytotoxic T cells or the like, which are necessary for T cell transfer therapy, can be induced *in vitro* in a short time. CTLs cannot be induced using dendritic cells that are not introduced with the minus-strand RNA virus. According to hitherto reported characteristics of other vectors, it is difficult to induce CTLs *in vitro* by gene transfer of such other vectors alone. Thus, the minus-strand RNA viral vector has 15 the advantage that it can activate T cells (CTL induction) merely by its introduction (see Fig. 19). In addition, the minus-strand RNA viral vector is superior to other vectors, due to the fact that it possesses all the characteristics of high introduction efficiency, stability of gene transfer, convenience, safety, and maintenance of the ability of the dendritic cells to activate T cells.

[0043]

20 Furthermore, the minus-strand RNA viral vector is also useful for differentiating stem cells after gene transfer into dendritic cells. When dendritic cell differentiation is induced after the minus-strand RNA viral vector is introduced into stem cells, the gene transfer efficiency reaches to nearly 70%. This efficiency is comparable to those of modified retroviral and lentiviral vectors. The introduction of adenoviral vector into stem cells is difficult, due to the 25 reduction in the expression level by the episome dilution after introduction. The minus-strand RNA virus can be applied for both methods wherein dendritic cell differentiation is performed after vector introduction into stem cells and wherein the gene is introduced into dendritic cells that have been differentiated from peripheral blood mononuclear cells.

[0044]

30 In addition, when the MOI is set high (for example, 10 or higher, preferably 20 or higher, more preferably 30 or higher, for example, 40 or higher, or 50 or higher), the minus-strand RNA viral vector can be stably introduced into cells with an introduction efficiency of nearly 100% without any significant influence on cell cytotoxicity. Moreover, since the minus-strand RNA virus does not integrate genes into the host genome, it has the advantage of a low risk of tumor 35 development.

[0045]

Herein, a minus-strand RNA virus refers to viruses that include a minus strand (an antisense strand corresponding to a sense strand encoding viral proteins) RNA as the genome. The minus-strand RNA is also referred to as negative strand RNA. The minus-strand RNA virus used in the present invention particularly includes single-stranded minus-strand RNA viruses (also referred to as non-segmented minus-strand RNA viruses). The “single-strand negative strand RNA virus” refers to viruses having a single-stranded negative strand [*i.e.*, a minus strand] RNA as the genome. Such viruses include viruses belonging to Paramyxoviridae (including the genera *Paramyxovirus*, *Morbillivirus*, *Rubulavirus*, and *Pneumovirus*), Rhabdoviridae (including the genera *Vesiculovirus*, *Lyssavirus*, and *Ephemerovirus*), Filoviridae, Orthomyxoviridae, (including Influenza viruses A, B, and C, and Thogoto-like viruses), Bunyaviridae (including the genera *Bunyavirus*, *Hantavirus*, *Nairovirus*, and *Phlebovirus*), Arenaviridae, and the like.

[0046]

In addition, the minus-strand RNA viral vector is a minus-strand RNA virus-based virion with infectivity and a vehicle for introducing genes into cells. Herein, “infectivity” refers to the capability of a minus-strand RNA viral vector to maintain cell-adhesion ability and introduce a gene carried by the vector to the inside of the cell to which the vector has adhered. In a preferable embodiment, the minus-strand RNA viral vector of this invention has a foreign gene incorporated into its genomic RNA for expression. The minus-strand RNA viral vector of this invention may have propagation ability or may be a defective-type vector with no propagation ability. “Having propagation ability” means that when a viral vector infects a host cell, the virus is replicated in the cell to produce infectious virions.

[0047]

“Recombinant virus” refers to a virus produced through a recombinant polynucleotide, or an amplification product thereof. “Recombinant polynucleotide” refers to a polynucleotide in which nucleotides are not linked at one or both ends as in the natural condition. Specifically, a recombinant polynucleotide is a polynucleotide in which the linkage of the polynucleotide chain has been artificially modified (cleaved and/or linked). Recombinant polynucleotides can be produced by using gene recombination methods known in the art in combination with polynucleotide synthesis, nuclease treatment, ligase treatment, etc. A recombinant virus can be produced by expressing a polynucleotide encoding a viral genome constructed through gene manipulation and reconstructing the virus. For example, methods for reconstructing a virus from cDNA that encodes the viral genome are known (Y. Nagai, A. Kato, *Microbiol. Immunol.*, 43, 613-624 (1999)).

[0048]

In the present invention, “gene” refers to a genetic substance, a nucleic acid coding a

transcriptional unit. Genes may be RNAs or DNAs. In this invention, a nucleic acid encoding a protein is referred to as a gene of that protein. Further, a gene may not encode a protein. For example, a gene may encode a functional RNA, such as a ribozyme or antisense RNA. A gene may be a naturally-occurring or artificially designed sequence. Furthermore, in the present
5 invention, "DNA" includes both single-stranded and double-stranded DNAs. Moreover, "encoding a protein" means that a polynucleotide includes an ORF that encodes an amino acid sequence of the protein in a sense or antisense strand, so that the protein can be expressed under appropriate conditions.

[0049]

10 A minus-strand RNA virus preferably used in the context of the present invention includes, for example, Sendai virus, Newcastle disease virus, Mumps virus, Measles virus, Respiratory syncytial virus (RS virus), rinderpest virus, distemper virus, simian parainfluenza virus (SV5), and human parainfluenza viruses 1, 2, and 3 belonging to *Paramyxoviridae*; Influenza virus belonging to *Orthomyxoviridae*; and Vesicular stomatitis virus and rabies virus
15 belonging to *Rhabdoviridae*.

[0050]

Further examples of virus that may be used in the context of the present invention include those selected from the group consisting of: Sendai virus (SeV), human parainfluenza virus-1 (HPIV-1), human parainfluenza virus-3 (HPIV-3), phocine distemper virus (PDV), canine
20 distemper virus (CDV), dolphin molbillivirus (DMV), peste-des-petits-ruminants virus (PDPR), measles virus (MV), rinderpest virus (RPV), Hendra virus (Hendra), Nipah virus (Nipah), human parainfluenza virus-2 (HPIV-2), simian parainfluenza virus 5 (SV5), human parainfluenza virus-4a (HPIV-4a), human parainfluenza virus-4b (HPIV-4b), mumps virus (Mumps), and Newcastle disease virus (NDV). A more preferred example is a virus selected from the group
25 consisting of Sendai virus (SeV), human parainfluenza virus-1 (HPIV-1), human parainfluenza virus-3 (HPIV-3), phocine distemper virus (PDV), canine distemper virus (CDV), dolphin molbillivirus (DMV), peste-des-petits-ruminants virus (PDPR), measles virus (MV), rinderpest virus (RPV), Hendra virus (Hendra), and Nipah virus (Nipah).

[0051]

30 More preferably, viruses of the present invention are preferably those belonging to *Paramyxoviridae* (including *Respirovirus*, *Rubulavirus*, and *Morbillivirus*) or derivatives thereof, and more preferably those belonging to the genus *Respirovirus* (also referred to as *Paramyxovirus*) or derivatives thereof. The derivatives include viruses that are genetically-modified or chemically-modified in a manner not to impair their gene-transferring
35 ability. Examples of viruses of the genus *Respirovirus* applicable to this invention are human parainfluenza virus-1 (HPIV-1), human parainfluenza virus-3 (HPIV-3), bovine parainfluenza

virus-3 (BPIV-3), Sendai virus (also referred to as murine parainfluenza virus-1), and simian parainfluenza virus-10 (SPIV-10). A more preferred paramyxovirus in this invention is the Sendai virus. These viruses may be derived from natural strains, wild strains, mutant strains, laboratory-passaged strains, artificially constructed strains, or the like.

5 [0052]

Genes harbored on a minus-strand RNA viral vector are situated in the antisense direction in the genomic RNA. Genomic RNA refers to RNA that has the function to form a ribonucleoprotein (RNP) with the viral proteins of a minus-strand RNA virus. Genes contained in the genome are expressed by the RNP, genomic RNA is replicated, and daughter RNPs are
10 formed. In general, the genome of a minus-strand RNA virus is constituted so that the viral genes are situated in an antisense orientation between the 3'-leader region and 5'-trailer region. Between the ORFs of individual genes exists a transcription ending sequence (E sequence) – intervening sequence (I sequence) – transcription starting sequence (S sequence) that allows the RNA encoding each ORF to be transcribed as a separate cistron.

15 [0053]

Genes encoding the viral proteins of a minus-strand RNA virus include NP, P, M, F, HN, and L genes. "NP, P, M, F, HN, and L genes" refer to genes encoding nucleocapside-, phospho-, matrix-, fusion-, hemagglutinin-neuraminidase-, and large-proteins respectively. Genes in each virus belonging to Paramyxovirinae are commonly listed as follows. In general,
20 NP gene is also listed as "N gene."

<i>Respirovirus</i>	NP	P/C/V	M	F	HN	-	L
<i>Rubulavirus</i>	NP	P/V	M	F	HN	(SH)	L
<i>Morbillivirus</i>	NP	P/C/V	M	F	H	-	L

For example, the database accession numbers for the nucleotide sequences of each of the
25 Sendai virus genes are: M29343, M30202, M30203, M30204, M51331, M55565, M69046, and X17218 for NP gene; M30202, M30203, M30204, M55565, M69046, X00583, X17007, and X17008 for P gene; D11446, K02742, M30202, M30203, M30204, M69046, U31956, X00584, and X53056 for M gene; D00152, D11446, D17334, D17335, M30202, M30203, M30204, M69046, X00152, and X02131 for F gene; D26475, M12397, M30202, M30203, M30204,
30 M69046, X00586, X02808, and X56131 for HN gene; and D00053, M30202, M30203, M30204, M69040, X00587, and X58886 for L gene. Examples of viral genes encoded by other viruses are: CDV, AF014953; DMV, X75961; HPIV-1, D01070; HPIV-2, M55320; HPIV-3, D10025; Mapuera, X85128; Mumps, D86172; MV, K01711; NDV, AF064091; PDPR, X74443; PDV, X75717; RPV, X68311; SeV, X00087; SV5, M81442; and Tupaia, AF079780 for N gene; CDV,
35 X51869; DMV, Z47758; HPIV-1, M74081; HPIV-3, X04721; HPIV-4a, M55975; HPIV-4b, M55976; Mumps, D86173; MV, M89920; NDV, M20302; PDV, X75960; RPV, X68311; SeV,

M30202; SV5, AF052755; and Tupaia, AF079780 for P gene; CDV, AF014953; DMV, Z47758; HPIV-1, M74081; HPIV-3, D00047; MV, AB016162; RPV, X68311; SeV, AB005796; and Tupaia, AF079780 for C gene; CDV, M12669; DMV, Z30087; HPIV-1, S38067; HPIV-2, M62734; HPIV-3, D00130; HPIV-4a, D10241; HPIV-4b, D10242; Mumps, D86171; MV, AB012948; NDV, AF089819; PDPR, Z47977; PDV, X75717; RPV, M34018; SeV, U31956; and SV5, M32248 for M gene; CDV, M21849; DMV, AJ224704; HPN-1, M22347; HPIV-2, M60182; HPIV-3, X05303; HPIV-4a, D49821; HPIV-4b, D49822; Mumps, D86169; MV, AB003178; NDV, AF048763; PDPR, Z37017; PDV, AJ224706; RPV, M21514; SeV, D17334; and SV5, AB021962 for F gene; and, CDV, AF112189; DMV, AJ224705; HPIV-1, U709498; HPIV-2, D000865; HPIV-3, AB012132; HPIV-4A, M34033; HPIV-4B, AB006954; Mumps, X99040; MV, K01711; NDV, AF204872; PDPR, Z81358; PDV, Z36979; RPV, AF132934; SeV, U06433; and SV-5, S76876 for HN (H or G) gene. However, a number of strains are known for each virus, and genes exist that include sequences other than those cited above, due to strain variation.

[0054]

The ORFs encoding these viral proteins and ORFs of the foreign genes are arranged in the antisense direction in the genomic RNAs, via the above-described E-I-S sequence. The ORF closest to the 3'-end of the genomic RNAs requires only an S sequence between the 3'-leader region and the ORF, and does not require an E or I sequence. Further, the ORF closest to the 5'-end of the genomic RNA requires only an E sequence between the 5'-trailer region and the ORF, and does not require an I or S sequence. Furthermore, two ORFs can be transcribed as a single cistron, for example, by using an internal ribosome entry site (IRES) sequence. In such a case, an E-I-S sequence is not required between these two ORFs. For example, in wild type paramyxoviruses, a typical RNA genome includes a 3'-leader region, six ORFs encoding the N, P, M, F, HN, and L proteins in the antisense direction and in this order, and a 5'-trailer region on the other end. The viral gene orientation in the genomic RNAs of the present invention is not restricted, but similarly to the wild type viruses, it is preferable that ORFs encoding the N, P, M, F, HN, and L proteins are arranged in this order, after the 3'-leader region, and before the 5'-trailer region. Certain types of viruses have different viral genes, but even in such cases, it is preferable that each gene be arranged as in the wild type, as described above. In general, vectors maintaining the N, P, and L genes can autonomously express genes from the RNA genome in cells, replicating the genomic RNA. Furthermore, by the action of genes such as the F and HN genes, which encode envelope proteins, and the M gene, infectious virions are formed and released to the outside of cells. Thus, such vectors become viral vectors with propagation ability. A foreign gene to be transduced into dendritic cells may be inserted into a protein-noncoding region in this genome, as described below.

[0055]

Further, a minus-strand RNA viral vector of this invention may be deficient in any of the wild type virus genes. For example, a viral vector that excludes the M, F, or HN gene, or any combination thereof, can be preferably used in this invention. Such viral vectors can be reconstituted, for example, by externally supplying the products of the deficient genes. Similar to wild type viruses, the viral vectors thus prepared adhere to host cells and cause cell fusion, but they cannot form daughter virions that retain the same infectivity as the original vector, because the vector genome introduced into cells is deficient in viral genes. Therefore, such vectors are useful as safe viral vectors that can only introduce genes once. Examples of genes in which the genome may be deficient are the F gene and/or HN gene. For example, viral vectors can be reconstituted by transfecting host cells with a plasmid expressing a recombinant minus-strand RNA viral vector genome deficient in the F gene, along with an F protein expression vector and expression vectors for the NP, P, and L proteins (WO00/70055 and WO00/70070; Li, H.-O. *et al.*, J. Virol. 74 (14) 6564-6569 (2000)). Viruses can also be produced, for example, using host cells that have incorporated the F gene into their chromosomes. In these proteins, the amino acid sequences do not need to be the same as the viral sequences, and a mutant or homologous gene from another virus may be used as a substitute, so long as the activity in nucleic acid introduction is the same as, or greater than, that of the natural type.

[0056]

Further, vectors that include an envelope protein other than that of the virus from which the vector genome was derived, may be prepared as viral vectors used in this invention. For example, when reconstituting a virus, a viral vector including a desired envelope protein can be generated by expressing an envelope protein other than the envelope protein encoded by the basic viral genome. Such proteins are not particularly limited. A desired protein that confers an ability to infect cells may be used. Examples of such proteins include the envelope proteins of other viruses, for example, the G protein of vesicular stomatitis virus (VSV-G). The VSV-G protein may be derived from an arbitrary VSV strain. For example, VSV-G proteins derived from Indiana serotype strains (J. Virology 39: 519-528 (1981)) may be used, but the present invention is not limited thereto. Furthermore, the present vector may include any arbitrary combination of envelope proteins derived from other viruses. Preferred examples of such proteins are envelope proteins derived from viruses that infect human cells. Such proteins are not particularly limited, and include retroviral amphotropic envelope proteins and the like. For example, the envelope proteins derived from mouse leukemia virus (MuLV) 4070A strain can be used as the retroviral amphotropic envelope proteins. In addition, envelope proteins derived from MuMLV 10A1 strain may also be used (for example, pCL-10A1 (Imgenex) (Naviaux, R. K. *et al.*, J. Virol. 70: 5701-5705 (1996)). The proteins of *Herpesviridae* include, for example, gB,

gD, gH, and gp85 proteins of herpes simplex viruses, and gp350 and gp220 proteins of EB virus. The proteins of *Hepadnaviridae* include the S protein of hepatitis B virus. These proteins may be used as fusion proteins in which the extracellular domain is linked to the intracellular domain of the F or HN protein. As described above, the viral vectors used in this invention include
5 pseudotype viral vectors that include envelope proteins, such as VSV-G, derived from viruses other than the virus from which the genome was derived. If the viral vectors are designed such that these envelope proteins are not encoded in RNA genomes, the proteins will never be expressed after virion infection of the cells.

[0057]

10 Furthermore, the viral vectors used in this invention may be, for example, vectors that include on the envelope surface thereof, proteins such as adhesion factors capable of adhering to specific cells, ligands, receptors, antibodies or fragments, or vectors that include a chimeric protein with these proteins in the extracellular domain and polypeptides derived from the virus envelope in the intracellular domain. Thus, the dendritic cell specificity of the vectors can be
15 controlled. These proteins may be encoded in the viral genome, or supplied through the expression of genes not in the viral genome (for example, genes carried by other expression vectors, or genes in the host chromosomes) at the time of viral vector reconstitution.

[0058]

Further, in the viral vectors, any viral gene contained in the vector may be modified
20 from the wild type gene in order to reduce the immunogenicity caused by viral proteins, or to enhance RNA transcriptional or replicational efficiency, for example. Specifically, for example, modifying at least one of the replication factors N, P, and L genes, is considered to enhance transcriptional or replicational function. Furthermore, although the HN protein, which is an envelope protein, has both hemagglutinin activity and neuraminidase activity, it is possible, for
25 example, to improve viral stability in blood if the former activity is attenuated, and infectivity can be controlled if the latter activity is modified. Further, it is also possible to control membrane fusion ability by modifying the F protein. For example, the epitopes of the F protein and/or HN protein, which can be cell surface antigenic molecules, can be analyzed, and using this, viral vectors with reduced antigenicity to these proteins can be prepared.

30 [0059]

Furthermore, the minus-strand RNA viral vector may be deficient in one or more accessory gene. For example, by knocking out the V gene, one of the SeV accessory genes, the pathogenicity of SeV toward hosts such as mice is remarkably reduced, without hindering gene expression and replication in cultured cells (Kato, A. *et al.*, 1997, J. Virol. 71: 7266-7272; Kato,
35 A. *et al.*, 1997, EMBO J. 16: 578-587; Curran, J. *et al.*, WO01/04272, EP1067179). Such attenuated vectors are particularly useful as nontoxic viral vectors for *in vivo* or *ex vivo* gene

transfer.

[0060]

Minus-strand RNA viruses are excellent gene transfer vectors. They do not have DNA phase and carry out transcription and replication only in the host cytoplasm, and consequently, chromosomal integration does not occur (Lamb, R.A. and Kolakofsky, D., Paramyxoviridae: The viruses and their replication. *In*: Fields BN, Knipe DM, Howley PM, (eds). Fields of Virology. Vol. 2. Lippincott - Raven Publishers: Philadelphia, 1996, pp. 1177-1204). Therefore, safety issues such as transformation and immortalization due to chromosomal aberration do not occur. This characteristic of minus-strand RNA viruses contributes greatly to safety when it is used as a vector. For example, results on foreign gene expression show that even after multiple continuous passages of SeV, almost no nucleotide mutation is observed. This suggests that the viral genome is highly stable and the inserted foreign genes are stably expressed over long periods of time (Yu, D. *et al.*, Genes Cells 2, 457-466 (1997)). Further, there are qualitative advantages associated with SeV not having a capsid structural protein, such as packaging flexibility and insert gene size, suggesting that minus-strand RNA viral vectors may become a novel class of highly efficient vectors for human gene therapy. SeV vectors with propagation ability are capable of introducing foreign genes of up to at least 4 kb in size, and can simultaneously express two or more kinds of genes by adding the transcriptional units.

[0061]

Further, Sendai virus is known to be pathogenic in rodents causing pneumonia, but is not pathogenic for human. This is also supported by a previous report that nasal administration of wild type Sendai virus does not have severely harmful effects on non-human primates (Hurwitz, J.L. *et al.*, Vaccine 15: 533-540, 1997). These Sendai virus characteristics suggest that Sendai virus vectors can be applied therapeutically on humans, supporting the proposition that Sendai virus vectors are a promising choice of gene therapy that targets human dendritic cells.

[0062]

Viral vectors are capable of encoding foreign genes in their genomic RNA. A recombinant viral vector harboring a foreign gene is obtained by inserting a foreign gene into an above-described viral vector genome. The foreign gene can be any desired gene that needs to be expressed in a target dendritic cell, and may be a gene that encodes a naturally-occurring protein, or protein modified from a naturally-occurring protein by deletion, substitution, or insertion. The foreign gene can be inserted at any desired position in a protein-noncoding region of the virus genome, for example. The above nucleic acid can be inserted, for example, between the 3'-leader region and the viral protein ORF closest to the 3'-end; between each of the viral protein ORFs; and/or between the viral protein ORF closest to the 5'-end and the 5'-trailer

region in genomic DNA. Further, in genomes deficient in the F or HN gene or the like, nucleic acids encoding the foreign genes can be inserted into those deficient regions. When introducing a foreign gene into a paramyxovirus, it is desirable to insert the gene such that the chain length of the polynucleotide to be inserted into the genome will be a multiple of six (Journal of Virology, Vol. 67, No. 8, 4822-4830, 1993). An E-I-S sequence should be arranged between the inserted foreign gene and the viral ORF. Two or more genes can be inserted in tandem via E-I-S sequences.

[0063]

Expression levels of a foreign gene carried in a vector can be controlled using the type of transcriptional initiation sequence added upstream (to the 3'-side of the negative strand) of the gene (WO01/18223). The expression levels can also be controlled by the position at which the foreign gene is inserted in the genome: the nearer to the 3'-end of the negative strand the insertion position is, the higher the expression level; while the nearer to the 5'-end the insertion position is, the lower the expression level. Thus, to obtain a desired gene expression level, the insertion position of a foreign gene can be appropriately controlled such that the combination with genes encoding the viral proteins before and after the foreign gene is most suitable. In general, since a high foreign gene expression level is thought to be advantageous, it is preferable to link the foreign gene to a highly efficient transcriptional initiation sequence, and to insert it near the 3'-end of the negative strand genome. Specifically, a foreign gene is inserted between the 3'-leader region and the viral protein ORF closest to the 3'-end. Alternatively, a foreign gene may be inserted between the ORFs of the viral gene closest to the 3'-end and the second closest viral gene. In wild type paramyxoviruses, the viral protein gene closest to the 3'-end of the genome is the N gene, and the second closest gene is the P gene. Alternatively, when a high level of expression of the introduced gene is undesirable, the gene expression level from the viral vector can be suppressed to obtain an appropriate effect, for example, by inserting the foreign gene at a site in the vector as close as possible to the 5'-side of the negative strand, or by selecting an inefficient transcriptional initiation sequence.

[0064]

To prepare a minus-strand RNA viral vector, a cDNA encoding a genomic RNA of a virus is transcribed in mammalian cells, in the presence of viral proteins (i.e., N, P, and L proteins) essential for reconstitution of an RNP, which is a component of a virus. Viral RNP can be reconstituted by producing either the negative strand genome (that is, the same antisense strand as the viral genome) or the positive strand (the sense strand encoding the viral proteins). Production of the positive strand is preferable for increased efficiency of vector reconstitution. The RNA terminals preferably reflect the terminals of the 3'-leader sequence and 5'-trailer sequence as accurately as possible, as in the natural viral genome. To accurately regulate the

5'-end of the transcript, for example, the RNA polymerase may be expressed within a cell using the recognition sequence of T7 RNA polymerase as a transcription initiation site. To regulate the 3'-end of the transcript, for example, a self-cleaving ribozyme can be encoded at the 3'-end of the transcript, allowing accurate cleavage of the 3'-end with this ribozyme (Hasan, M. K. *et al.*, J. Gen. Virol. 78: 2813-2820, 1997; Kato, A. *et al.*, 1997, EMBO J. 16: 578-587; and Yu, D. *et al.*, 1997, Genes Cells 2: 457-466).

[0065]

For example, a recombinant Sendai virus vector carrying a foreign gene can be constructed as follows, according to descriptions in: Hasan, M. K. *et al.*, J. Gen. Virol. 78: 2813-2820, 1997; Kato, A. *et al.*, 1997, EMBO J. 16: 578-587; Yu, D. *et al.*, 1997, Genes Cells 2: 457-466; or the like.

First, a DNA sample including a cDNA sequence of an objective foreign gene is prepared. The DNA sample is preferably one that can be confirmed to be a single plasmid by electrophoresis at a concentration of 25 ng/ μ l or more. The following explains the case of using a *NotI* site to insert a foreign gene into a DNA encoding a viral genomic RNA, with reference to examples. When a *NotI* recognition site is included in a target cDNA nucleotide sequence, the base sequence is altered using site-directed mutagenesis or the like, such that the encoded amino acid sequence does not change, and the *NotI* site is preferably excised in advance. The objective gene fragment is amplified from this sample by PCR, and then recovered. By adding the *NotI* site to the 5' regions of a pair of primers, both ends of the amplified fragments become *NotI* sites. E-I-S sequences are designed to be included in primers such that, after a foreign gene is inserted into the viral genome, one E-I-S sequence each is placed between the ORF of the foreign gene, and either side of the ORFs of the viral genes.

[0066]

For example, to guarantee cleavage with *NotI*, the forward side synthetic DNA sequence has a form in which any desired sequence of not less than two nucleotides (preferably four nucleotides not including a sequence derived from the *NotI* recognition site, such as GCG and GCC, and more preferably ACTT) is selected at the 5'-side, and a *NotI* recognition site 'gcggccgc' is added to its 3'-side. To that 3'-side, nine arbitrary nucleotides, or nine plus a multiple of six nucleotides are further added as a spacer sequence. To the further 3' of this, a sequence corresponding to about 25 nucleotides of the ORF of a desired cDNA, including and counted from the initiation codon ATG, is added. The 3'-end of the forward side synthetic oligo DNA is preferably about 25 nucleotides, selected from the desired cDNA such that the final nucleotide becomes a G or C.

[0067]

For the reverse side synthetic DNA sequence, no less than two arbitrary nucleotides

(preferably four nucleotides not including a sequence derived from a *NotI* recognition site, such as GCG and GCC, and more preferably ACTT) are selected from the 5'-side, a *NotI* recognition site 'gcggccgc' is added to its 3'-side, and to that 3' is further added an oligo DNA insert fragment for adjusting the length. The length of this oligo DNA is designed such that the chain length of the *NotI* fragment of the final PCR-amplified product will become a multiple of six nucleotides (the so-called "rule of six"); Kolakofski, D., *et al.*, J. Virol. 72: 891-899, 1998; Calain, P. and Roux, L., J. Virol. 67: 4822-4830, 1993; Calain, P. and Roux, L., J. Virol. 67: 4822-4830, 1993). When adding an E-I-S sequence to this primer, to the 3'-side of the oligo DNA insertion fragment is added the complementary strand sequence of the Sendai virus S, I, and E sequences, preferably 5'-CTTTCACCCT-3' (SEQ ID NO: 1), 5'-AAG-3', and 5'-TTTTTCTTACTACGG-3' (SEQ ID NO: 2), respectively; and further to this 3'-side is added a complementary strand sequence corresponding to about 25 nucleotides, counted backwards from the termination codon of a desired cDNA sequence, whose length has been selected such that the final nucleotide of the chain becomes a G or C, to make the 3'-end of the reverse side synthetic DNA.

[0068]

PCR can be performed according to conventional methods, using Taq polymerase or other DNA polymerases. Objective amplified fragments may be digested with *NotI*, and then inserted into the *NotI* site of plasmid vectors such as pBluescript. The nucleotide sequences of PCR products thus obtained are confirmed with a sequencer, and plasmids that include the correct sequence are selected. The inserted fragment is excised from these plasmids using *NotI*, and cloned into the *NotI* site of a plasmid composed of genomic cDNA. A recombinant Sendai virus cDNA can also be obtained by inserting the fragment directly into the *NotI* site of a genomic cDNA, without using a plasmid vector.

[0069]

For example, a recombinant Sendai virus genomic cDNA can be constructed according to methods described in the literature (Yu, D. *et al.*, Genes Cells 2: 457-466, 1997; Hasan, M. K. *et al.*, J. Gen. Virol. 78: 2813-2820, 1997). For example, an 18 bp spacer sequence (5'-(G)-CGGCCGCAGATCTTCACG-3') (SEQ ID NO: 3), including a *NotI* restriction site, is inserted between the leader sequence and the ORF of N protein of the cloned Sendai virus genomic cDNA (pSeV(+)), obtaining plasmid pSeV18⁺b(+), which includes an auto-cleavage ribozyme site derived from the antigenomic strand of delta hepatitis virus (Hasan, M. K. *et al.*, 1997, J. General Virology 78: 2813-2820). A recombinant Sendai virus cDNA including a desired foreign gene can be obtained by inserting a foreign gene fragment into the *NotI* site of pSeV18⁺b(+).

[0070]

A viral vector can be reconstituted by transcribing a DNA encoding a genomic RNA of a recombinant virus thus prepared, in cells in the presence of the above-described viral proteins (L, P, and N). The present invention provides minus-strand RNA viral vectors for transfer into dendritic cells. In addition, the present invention relates to the use of the minus-strand RNA viral vectors in the preparation of dendritic cells introduced with a gene and in the preparation of mature dendritic cells. The present invention also provides DNAs encoding the viral genomic RNAs of the minus-strand RNA viral vectors for manufacturing the minus-strand RNA viral vectors for transfer into dendritic cells. This invention also relates to the use of DNAs encoding the genomic RNAs of the vectors, in the manufacture of the vectors of this invention. The recombinant viruses can be reconstituted by methods known in the art (WO97/16539; WO97/16538; Durbin, A. P. *et al.*, 1997, *Virology* 235: 323-332; Whelan, S. P. *et al.*, 1995, *Proc. Natl. Acad. Sci. USA* 92: 8388-8392; Schnell, M. J. *et al.*, 1994, *EMBO J.* 13: 4195-4203; Radecke, F. *et al.*, 1995, *EMBO J.* 14: 5773-5784; Lawson, N. D. *et al.*, *Proc. Natl. Acad. Sci. USA* 92: 4477-4481; Garcin, D. *et al.*, 1995, *EMBO J.* 14: 6087-6094; Kato, A. *et al.*, 1996, *Genes Cells* 1: 569-579; Baron, M. D. and Barrett, T., 1997, *J. Virol.* 71: 1265-1271; Bridgen, A. and Elliott, R. M., 1996, *Proc. Natl. Acad. Sci. USA* 93: 15400-15404). With these methods, minus strand RNA viruses including parainfluenza virus, vesicular stomatitis virus, rabies virus, measles virus, rinderpest virus, and Sendai virus can be reconstituted from DNA. The vectors of this invention can be reconstituted according to these methods. When a viral vector DNA is made F gene, HN gene, and/or M gene deficient, such DNAs do not form infectious virions as is. However, infectious virions can be formed by separately introducing host cells with these deficient genes, and/or genes encoding the envelope proteins of other viruses, and then expressing these genes therein.

[0071]

Specifically, the viruses can be prepared by the steps of: (a) transcribing cDNAs encoding genomic RNAs of minus-strand RNA viruses (negative strand RNAs), or complementary strands thereof (positive strands), in cells expressing N, P, and L proteins; and (b) harvesting culture supernatants thereof including the produced minus-strand RNA viruses. For transcription, a DNA encoding a genomic RNA is linked downstream of an appropriate promoter. The genomic RNA thus transcribed is replicated in the presence of N, L, and P proteins to form an RNP complex. Then, in the presence of M, HN, and F proteins, virions enclosed in an envelope are formed. For example, a DNA encoding a genomic RNA can be linked downstream of a T7 promoter, and transcribed to RNA by T7 RNA polymerase. Any desired promoter can be used as a promoter, in addition to those including a T7 polymerase recognition sequence. Alternatively, RNA transcribed *in vitro* may be transfected into cells.

[0072]

Enzymes essential for the initial transcription of genomic RNA from DNA, such as T7 RNA polymerase, can be supplied by transducing the plasmid or viral vectors that express them, or, for example, by incorporating the RNA polymerase gene into a chromosome of the cell so as to enable induction of its expression, and then inducing expression at the time of viral reconstitution. Further, genomic RNA and viral proteins essential for vector reconstitution are supplied, for example, by transducing the plasmids that express them. In supplying these viral proteins, helper viruses such as the wild type or certain types of mutant minus-strand RNA viruses are used.

[0073]

Methods for transducing DNAs expressing the genomic RNAs into cells include, for example, (i) methods for making DNA precipitates which target cells can internalize; (ii) methods for making complexes including DNAs that are suitable for internalization by target cells, and have a low-cytotoxic positive charge; and (iii) methods for using electric pulses to instantaneously create holes in the target cell membrane, which are of sufficient size for DNA molecules to pass through.

[0074]

In the context of method (ii), various transfection reagents can be used. For example, DOTMA (Roche), Superfect (QIAGEN #301305), DOTAP, DOPE, DOSPER (Roche #1811169), and the like can be cited. Regarding method (i), for example, transfection methods using calcium phosphate can be cited, and although DNAs transferred into cells by this method are internalized by phagosomes, a sufficient amount of DNA is known to enter the nucleus (Graham, F. L. and Van Der Eb, J., 1973, *Virology* 52: 456; Wigler, M. and Silverstein, S., 1977, *Cell* 11: 223). Chen and Okayama investigated the optimization of transfer techniques, reporting that (1) incubation conditions for cells and coprecipitates are 2 to 4% CO₂, 35°C, and 15 to 24 hours, (2) the activity of circular DNA is higher than linear DNA, and (3) optimal precipitation is obtained when the DNA concentration in the precipitate mixture is 20 to 30 µg/ml (Chen, C. and Okayama, H., 1987, *Mol. Cell. Biol.* 7: 2745). The methods of (ii) are suitable for transient transfections. Methods for performing transfection by preparing a DEAE-dextran (Sigma #D-9885 M.W. 5 x 10⁵) mixture with a desired DNA concentration ratio have been known for a while. Since most complexes are decomposed in endosomes, chloroquine may also be added to enhance the effect (Calos, M. P., 1983, *Proc. Natl. Acad. Sci. USA* 80: 3015). The methods of (iii) are referred to as electroporation methods, and are used more in general than methods (i) or (ii) because they are not cell-selective. The efficiency of these methods is presumed to be good under optimal conditions for: the duration of pulse electric current, shape of the pulse, potency of electric field (gap between electrodes, voltage), conductivity of buffer, DNA concentration, and cell density.

[0075]

Of the above three categories, the methods of (ii) are simple to operate and facilitate examination of many samples using a large amount of cells, making transfection reagents suitable for the transduction into cells of DNA for vector reconstitution. Preferably, the
5 Superfect Transfection Reagent (QIAGEN, Cat No. 301305), or the DOSPER Liposomal Transfection Reagent (Roche, Cat No. 1811169) is used; however, the transfection reagents are not limited to these.

[0076]

Specifically, virus reconstitution from cDNA can be carried out, for example, as
10 follows:

In a plastic plate of about 6 to 24 wells, or a 100-mm Petri dish or the like, simian kidney-derived LLC-MK2 cells (ATCC CCL-7) are cultured up to about 100% confluency, using minimum essential medium (MEM) including 10% fetal calf serum (FCS) and antibiotics (100 units/ml penicillin G and 100 µg/ml streptomycin). Then they are infected with, for example,
15 two plaque forming units (PFU)/cell of the recombinant vaccinia virus vTF7-3, which expresses T7 RNA polymerase and has been inactivated by 20-minutes of UV irradiation in the presence of 1 µg/ml psoralen (Fuerst, T. R. *et al.*, Proc. Natl. Acad. Sci. USA 83: 8122-8126, 1986; Kato, A. *et al.*, Genes Cells 1: 569-579, 1996). The amount of psoralen added and the UV irradiation time can be appropriately adjusted. One hour after infection, 2 to 60 µg, and more preferably 3
20 to 20 µg, of DNA encoding the genomic RNA of a recombinant Sendai virus is transfected along with the plasmids expressing trans-acting viral proteins essential for viral RNP production (0.5 to 24 µg of pGEM-N, 0.25 to 12 µg of pGEM-P, and 0.5 to 24 µg of pGEM-L) (Kato, A. *et al.*, Genes Cells 1: 569-579, 1996), using the lipofection method or the like with Superfect (QIAGEN). For example, the ratio of the amounts of expression vectors encoding the N, P, and
25 L proteins is preferably 2:1:2, and the plasmid amounts are appropriately adjusted in the range of 1 to 4 µg of pGEM-N, 0.5 to 2 µg of pGEM-P, and 1 to 4 µg of pGEM-L.

[0077]

The transfected cells are cultured, as desired, in serum-free MEM composed of 100 µg/ml of rifampicin (Sigma) and cytosine arabinoside (AraC), more preferably only 40 µg/ml of
30 cytosine arabinoside (AraC) (Sigma). Optimal drug concentrations are set so as to minimize cytotoxicity due to the vaccinia virus, and to maximize virus recovery rate (Kato, A. *et al.*, 1996, Genes Cells 1: 569-579). After culturing for about 48 to 72 hours after transfection, cells are harvested, and then disintegrated by repeating freeze-thawing three times. LLC-MK2 cells are re-infected with the disintegrated materials including RNP, and cultured. Alternatively, the
35 culture supernatant is recovered, added to a culture solution of LLC-MK2 cells to infect them, and the cells are then cultured. Transfection can be conducted by, for example, forming a

complex with lipofectamine, polycationic liposome, or the like, and transducing the complex into cells. Specifically, various transfection reagents can be used. For example, DOTMA (Roche), Superfect (QIAGEN #301305), DOTAP, DOPE, and DOSPER (Roche #1811169) may be cited. In order to prevent decomposition in the endosome, chloroquine may also be added (Calos, M. P.,
5 1983, Proc. Natl. Acad. Sci. USA 80: 3015). In cells transduced with RNP, viral gene expression from RNP and RNP replication progress, and the vector is amplified. By diluting the viral solution thus obtained (for example, 10^6 -fold), and then repeating the amplification, the vaccinia virus vTF7-3 can be completely eliminated. Amplification is repeated, for example, three or more times. Vectors thus obtained can be stored at -80°C . In order to reconstitute a
10 viral vector having no propagation ability and lacking a gene encoding an envelope protein, LLC-MK2 cells expressing the envelope protein may be used for transfection, or a plasmid expressing the envelope protein may be cotransfected. Alternatively, a defective type viral vector can be amplified by culturing the transfected cells overlaid with LLK-MK2 cells expressing the envelope protein (see WO00/70055 and WO00/70070).

15 [0078]

Titers of viruses thus recovered can be determined, for example, by measuring CIU (Cell-Infected Unit) or hemagglutination activity (HA) (WO00/70070; Kato, A. *et al.*, 1996, Genes Cells 1: 569-579; Yonemitsu, Y. & Kaneda, Y., Hemagglutinating virus of Japan-liposome-mediated gene delivery to vascular cells. Ed. by Baker AH. Molecular Biology
20 of Vascular Diseases. Method in Molecular Medicine: Humana Press: pp. 295-306, 1999). Titers of vectors carrying GFP (green fluorescent protein) marker genes and the like can be quantified by directly counting infected cells, using the marker as an indicator (for example, as GFP-CIU). Titers thus measured can be treated in the same way as CIU (WO00/70070).

[0079]

25 So long as a viral vector can be reconstituted, the host cells used in the reconstitution are not particularly limited. For example, in the reconstitution of Sendai virus vectors and the like, cultured cells such as LLC-MK2 cells and CV-1 cells derived from monkey kidney, BHK cells derived from hamster kidney, and cells derived from humans can be used. By expressing suitable envelope proteins in these cells, infectious virions including the envelope can also be
30 obtained. Further, to obtain a large quantity of a Sendai virus vector, a viral vector obtained from an above-described host can be infected to embrionated hen eggs, to amplify the vector. Methods for manufacturing viral vectors using hen eggs have already been developed (Nakanishi, *et al.*, ed. (1993), "State-of-the-Art Technology Protocol in Neuroscience Research III, Molecular Neuron Physiology", Koseisha, Osaka, pp. 153-172). Specifically, for example, a
35 fertilized egg is placed in an incubator, and cultured for nine to twelve days at 37 to 38°C to grow an embryo. After the viral vector is inoculated into the allantoic cavity, the egg is

cultured for several days (for example, three days) to proliferate the viral vector. Conditions such as the period of culture may vary depending upon the recombinant Sendai virus being used. Then, allantoic fluids including the vector are recovered. Separation and purification of a Sendai virus vector from allantoic fluids can be performed according to a usual method (Tashiro, M., "Virus Experiment Protocol," Nagai, Ishihama, ed., Medical View Co., Ltd., pp. 68-73, (1995)).

[0080]

For example, the construction and preparation of Sendai virus vectors defective in F gene can be performed as described below (see WO00/70055 and WO00/70070).

10 <1> Construction of a genomic cDNA of an F-gene defective Sendai virus, and a plasmid expressing F gene:

A full-length genomic cDNA of Sendai virus (SeV), the cDNA of pSeV18⁺ b (+) (Hasan, M. K. *et al.*, 1997, J. General Virology 78: 2813-2820) ("pSeV18⁺ b (+)" is also referred to as "pSeV18⁺"), is digested with *SphI/KpnI* to recover a fragment (14,673 bp), which is cloned into pUC18 to prepare plasmid pUC18/KS. Construction of an F gene-defective site is performed on this pUC18/KS. An F gene deficiency is created by a combination of PCR-ligation methods, and, as a result, the F gene ORF (ATG-TGA = 1,698 bp) is removed. Then, for example, 'atgcatgccggcagatga (SEQ ID NO: 4)' is ligated to construct an F gene-defective type SeV genomic cDNA (pSeV18⁺/ΔF). A PCR product formed in PCR by using the pair of primers [forward: 5'-gttgagtactgcaagagc/SEQ ID NO: 5, reverse: 5'-tttgcggcatgcatgtttccaaggggagagttttgcaacc/SEQ ID NO: 6] is connected upstream of F, and a PCR product formed using the pair of primers [forward: 5'-atgcatgccggcagatga/SEQ ID NO: 7, reverse: 5'-tgggtgaatgagagaatcagc/SEQ ID NO: 8] is connected downstream of the F gene with *EcoT22I*. The plasmid thus obtained is digested with *SacI* and *SalI* to recover a 4,931 bp fragment of the region including the F gene-defective site, which is cloned into pUC18 to form pUC18/dFSS. This pUC18/dFSS is digested with *DraIII*, the fragment is recovered, replaced with the *DraIII* fragment of the region including the F gene of pSeV18⁺, and ligated to obtain the plasmid pSeV18⁺/ΔF.

A foreign gene is inserted, for example, into the *NsiI* and *NgoMIV* restriction enzyme sites in the F gene-defective site of pUC18/dFSS. For this, a foreign gene fragment may be, for example, amplified using an *NsiI*-tailed primer and an *NgoMIV*-tailed primer.

[0081]

<2> Preparation of helper cells that induce SeV-F protein expression:

To construct an expression plasmid of the Cre/loxP induction type that expresses the Sendai virus F gene (SeV-F), the SeV-F gene is amplified by PCR, and inserted to the unique *SwaI* site of the plasmid pCALNdlw (Arai, T. *et al.*, J. Virology 72, 1998, p1115-1121), which is

designed to enable the inducible expression of a gene product by Cre DNA recombinase, thus constructing the plasmid pCALNdLw/F.

To recover infectious virions from the F gene-defective genome, a helper cell line expressing SeV-F protein is established. The monkey kidney-derived LLC-MK2 cell line, which is commonly used for SeV proliferation, can be used as the cells, for example. LLC-MK2 cells are cultured in MEM supplemented with 10% heat-inactivated fetal bovine serum (FBS), penicillin G sodium (50 units/ml), and streptomycin (50 µg/ml) at 37°C in 5% CO₂. Since the SeV-F gene product is cytotoxic, the above-described plasmid pCALNdLw/F, which was designed to enable inducible expression of the F gene product with Cre DNA recombinase, is transfected to LLC-MK2 cells by the calcium phosphate method (using a mammalian transfection kit (Stratagene)), according to protocols well known in the art.

The plasmid pCALNdLw/F (10 µg) is transduced into LLC-MK2 cells grown to 40% confluency using a 10-cm plate, and the cells are then cultured in MEM (10 ml) including 10% FBS, in a 5% CO₂ incubator at 37°C for 24 hours. After 24 hours, the cells are detached and suspended in the medium (10 ml). The suspension is then seeded into five 10-cm dishes, 5 ml into one dish, 2 ml each into two dishes, and 0.2 ml each into two dishes, and cultured in MEM (10 ml) including G418 (GIBCO-BRL) (1200 µg/ml) and 10% FBS. The cells were cultured for 14 days, exchanging the medium every two days, to select cell lines stably transduced with the gene. The cells grown from the above medium that show G418 resistance are recovered using a cloning ring. Culture of each clone thus recovered is continued in 10-cm plates until confluent.

After the cells have grown to confluency in a 6-cm dish, F protein expression can be induced by infecting the cells with adenovirus AxCANCre, for example, at MOI = 3, according to the method of Saito, *et al.* (Saito *et al.*, Nucl. Acids Res. 23: 3816-3821 (1995); Arai, T. *et al.*, J. Virol 72, 1115-1121 (1998)).

[0082]

<3> Reconstruction and amplification of F gene-deficient SeV virus:

The above-described plasmid pSeV18⁺/ΔF inserted with the foreign gene is transfected into LLC-MK2 cells by the procedure described below. LLC-MK2 cells are seeded on 100-mm dishes at 5 x 10⁶ cells/dish. To transcribe the genomic RNA using T7 RNA polymerase, the cells are cultured for 24 hours, and then recombinant vaccinia virus, which expresses T7 RNA polymerase (PLWUV-VacT7: Fuerst, T.R. *et al.*, Proc. Natl. Acad. Sci. USA 83, 8122-8126 (1986)) and is treated with psoralen and long-wavelength ultraviolet light (365 nm) for 20 minutes, is inoculated to the cells at a MOI of about 2 at room temperature for one hour. The ultraviolet light irradiation to the vaccinia virus can be achieved, for example, by using UV Stratalinker 2400 with five 15-watt bulbs (Catalog No. 400676 (100V); Stratagene, La Jolla, CA,

USA). After the cells are washed with serum-free MEM, plasmid expressing the genomic RNA and expression plasmids each expressing N, P, L, F, or HN protein of the minus-strand RNA virus are transfected into the cells using an appropriate lipofection reagent. The plasmid ratio is preferably, but is not limited to, 6:2:1:2:2 in this order. For example, the expression plasmid
5 for the genomic RNA, and the expression plasmids each of which expresses N, P, or L protein, or F and HN proteins (pGEM/NP, pGEM/P, pGEM/L, and pGEM/F-HN; WO00/70070, Kato, A. *et al.*, Genes Cells 1, 569-579 (1996)) are transfected at amounts of 12, 4, 2, 4, and 4 $\mu\text{g}/\text{dish}$, respectively. After a few hours of culture, the cells are washed twice with serum-free MEM, and then cultured in MEM supplemented with 40 $\mu\text{g}/\text{ml}$ cytosine β -D-arabinofuranoside (AraC:
10 Sigma, St. Louis, MO) and 7.5 $\mu\text{g}/\text{ml}$ trypsin (Gibco-BRL, Rockville, MD). The cells are recovered, and the resulting pellet is suspended in OptiMEM (10^7 cells/ml). The suspension is subjected to three freeze-thaw cycles, and mixed with lipofection reagent DOSPER (Boehringer Mannheim) (10^6 cells/25 μl DOSPER). After the mixture is allowed to stand at room temperature for 15 minutes, it is transfected to F-expressing helper cells (10^6 cells/well in
15 12-well-plate) cloned as described above. The cells are cultured in serum-free MEM (containing 40 $\mu\text{g}/\text{ml}$ AraC and 7.5 $\mu\text{g}/\text{ml}$ trypsin), and the supernatant is collected. Viruses deficient in genes other than F, for example, HN and M genes, can be prepared by a similar method as described above.

[0083]

20 For the preparation of viral gene-deficient vectors, for example, two or more types of vectors which differ in the deficient viral gene on the viral genome carried by the vector are introduced into same cells. The respective deficient viral proteins are supplied through the expression from the other vector(s). Therefore, the vectors complement each other to form infectious viral particles, resulting in a complete replication cycle and amplification of the viral
25 vectors. Specifically, when two or more types of vectors of the present invention are inoculated in combination that allows complementation of viral proteins, a mixture of the viral gene-deficient vectors can be produced on a large scale at low cost. Since such viruses lack viral genes, their genome sizes are smaller than those of viral gene-nondeficient viruses and thus can carry larger foreign genes. Furthermore, these viruses that are non-proliferative due to the
30 lack of viral genes become diluted outside cells, which make it difficult to maintain their coinfection. The viruses become sterile and thus are advantageous from the viewpoint of controlling their environmental release.

[0084]

35 There is no limitation on the foreign gene to be introduced using the minus-strand RNA virus, and naturally occurring proteins include, for example, hormones, cytokines, growth factors, receptors, intracellular signaling molecules, enzymes, and peptides. The proteins may be

secretory proteins, membrane proteins, cytoplasmic proteins, nuclear proteins, and the like. Artificial proteins include, for example, fusion proteins such as chimeric toxin, dominant negative proteins (including soluble receptor molecules or membrane bound dominant negative receptors), truncated cell adhesion molecules, and cell surface molecules. The proteins may also be proteins to which a secretory signal, membrane-localization signal, nuclear translocation signal, or the like has been attached. Functions of a particular gene can be suppressed by introducing and expressing antisense RNA molecule, RNA-cleaving ribozyme, or the like as the transfer gene. When a viral vector is prepared using a gene for treating diseases as the foreign gene, gene therapy can be performed through the introduction of the vector. The viral vector of the present invention is applicable to gene therapy wherein the genes are expressed by direct administration or by *ex vivo* administration, and enables expression of foreign genes for which therapeutic effect can be expected, internal genes short in *in vivo* supply, or the like. In addition, the method of the present invention can also be used as a gene therapy vector in regeneration medicine.

[0085]

According to the method for producing viruses as described herein, the viral vector of the present invention can be released into extracellular fluid of virus producing cells at a titer of, for example, 1×10^5 CIU/ml or higher, preferably 1×10^6 CIU/ml or higher, more preferably 5×10^6 CIU/ml or higher, more preferably 1×10^7 CIU/ml or higher, more preferably 5×10^7 CIU/ml or higher, more preferably 1×10^8 CIU/ml or higher, and more preferably 5×10^8 CIU/ml or higher. The titer of virus can be determined according to methods described herein or elsewhere (Kiyotani, K. *et al.*, *Virology* 177 (1), 65-74 (1990); and WO00/70070).

[0086]

The recovered viral vectors can be purified to be substantial pure. The purification can be achieved using known purification/separation methods, including filtration, centrifugation, adsorption, and column purification, or any combinations thereof. The phrase "substantially pure" means that the vector component constitutes a major proportion of a solution of the vector. For example, a viral vector composition can be confirmed to be substantially pure by the fact that the proportion of protein contained as the viral vector component to the total protein (excluding proteins added as carriers and stabilizers) in the solution is 10% (w/w) or greater, preferably 20% or greater, more preferably 50% or greater, preferably 70% or greater, more preferably 80% or greater, and even more preferably 90% or greater. Specific purification methods for, for example, the paramyxovirus vector includes methods using cellulose sulfate ester or cross-linked polysaccharide sulfate ester (Japanese Patent Application Kokoku Publication No. (JP-B) S62-30752 (examined, approved Japanese patent application published for opposition), JP-B S62-33879, and JP-B S62-30753) and methods including adsorbing to fucose sulfate-containing

polysaccharide and/or degradation products thereof (WO97/32010), but are not limited thereto.

[0087]

In the production of compositions containing the vector, the vector may be combined with desired pharmaceutically acceptable carriers or media according to needs. The

5 “pharmaceutically acceptable carriers or media” refers to materials that can be administered together with the vector and that do not significantly inhibit the gene transfer via the vector. Such carriers and media include, for example, sterile water, sodium chloride solution, dextrose solution, Ringer’s solution containing dextrose, sodium chloride, and lactated, culture medium, serum, and phosphate buffered saline (PBS). They may be appropriately combined with the

10 vector to formulate a composition. The composition of the present invention may also include carriers or media such as deionized water and dextrose solution. The composition may also include membrane stabilizers for liposome (for example, sterols such as cholesterol). The composition may also include antioxidants (for example, tocopherol or vitamin E). In addition, the composition may also include vegetable oils, suspending agents, detergents, stabilizers,

15 biocidal agents, and the like. Furthermore, preservatives and other additives may also be added. The formula of the present composition may be aqueous solution, capsule, suspension, syrup, or the like. The composition of the present invention may also be in a form of solution, freeze-dried product, or aerosol. When it is a freeze-dried product, it may include sorbitol, sucrose, amino acids, various proteins, and the like as a stabilizer. The composition containing

20 the vector of the present invention is useful as a reagent for introducing genes into dendritic cells and also as a pharmaceutical that is used in gene therapy targeting dendritic cells. Furthermore, the vector solution is useful as a vaccine (J. I. Mayordomo *et al.*, Nature Med. 1 (12), 1279-1302, (1995)). Moreover, when an antigen peptide is expressed in dendritic cells using the vector of the present invention, the cells presenting this peptide can be used as a vaccine. The vaccine

25 compositions may include immunostimulants, such as cytokine, cholera toxin, and Salmonella toxin to improve immunogenicity. Furthermore, the vaccine may be combined with adjuvants, such as alum, incomplete Freund’s adjuvant, MF59 (oil emulsion), MTP-PE (muramyl tripeptide derived from cell wall of mycobacteria), and QS-21 (derived from soapbark tree *Quilaja saponaria*).

[0088]

When administering the composition, it is effective to combine them with cytokines that improve the adjuvant effect. Such genes include, for example, (i) a combination of IL-2 and single-chain IL-12 (Proc. Natl. Acad. Sci. USA 96 (15): 8591-8596, 1999); (ii) IL-2 and interferon- γ (U.S. Patent No. 5,798,100); (iii) granulocyte colony stimulating factor (GM-CSF),

35 which is used alone; and (iv) a combination of GM-CSF and IL-4 (J. Neurosurgery 90 (6), 1115-1124 (1999)).

[0089]

An antigen to be presented by dendritic cells may be encoded by the minus-strand RNA viral vector, added to (specifically, pulsed into) dendritic cells into which the vector has been introduced, or expressed using an alternate desired vector. Such antigens include desired
5 antigens related to infectious microorganisms, viruses, parasites, pathogens, cancers, and the like. These may be structural or non-structural proteins. Such antigens (or processed peptides thereof) are presented on the cell surface bound to MHC molecules on the surface of dendritic cells to induce immune responses.

[0090]

10 When used as a vaccine, the antigens can be applied to, for example, tumors, infectious diseases, and other general diseases. To treat infectious diseases, for example, epitopes of an antigen protein of an infectious microorganism may be analyzed, and then expressed or presented by dendritic cells.

Antigens derived from pathogens include, for example, proteins of hepatitis A virus,
15 hepatitis B virus, hepatitis C virus, hepatitis delta virus, papilloma virus antigen, herpes simplex virus (HSV), varicella-zoster virus (VZV), Epstein-Barr virus, Cytomegalovirus (CMV), HIV, malaria, and the like, or partial peptides thereof. The minus-strand RNA viruses encoding such antigen proteins can be used prophylactically or therapeutically. Specifically, envelopes of influenza highly-virulent strain H5N1 for influenza, envelope proteins (Vaccine, vol. 17, No.
20 15-16, 1869-1882 (1999)) for Japanese encephalitis, HIV and SIV gag proteins (J. Immunology (2000) vol. 164, 4968-4978), HIV envelope proteins, Nef protein, and other viral proteins for AIDS can be mentioned. In addition, for example, cholera toxin B subunit (CTB) (Arakawa T, *et al.*, Nature Biotechnology (1998) 16 (10): 934-8, Arakawa T, *et al.*, Nature Biotechnology (1998) 16 (3): 292-7) for cholera; rabies virus glycoprotein (Lodmell DL *et al.*, 1998, Nature
25 Medicine 4 (8):949-52) for rabies; and capsid protein L1 of human papilloma virus type 6 (J. Med. Virol, 60, 200-204 (2000)) for cervical carcinoma can be mentioned. Antigen proteins of other pathogenic viruses can also be expressed from the vector. Furthermore, it is possible to use JE-E antigen protein of Japanese encephalitis (Japanese Patent Application Kokai Publication No. (JP-A) S64-74982 (unexamined, published Japanese patent application), JP-A
30 H01-285498), gD2 protein of human herpes simplex virus (JP-A H05-252965), polypeptides derived from hepatitis C virus (JP-A H05-192160), polypeptides derived from pseudorabies virus (Japanese Patent Kohyo Publication No. (JP-A) H07-502173 (unexamined Japanese national phase publication corresponding to a non-Japanese international publication)), and the like. For example, cells derived from patients infected with such pathogenic microorganisms may be
35 analyzed to identify an epitope of an antigen protein to be presented on antigen-presenting cells (APC) for use. It is preferred to appropriately select the HLA type and identify an epitope

corresponding to the desired HLA for use.

[0091]

To specifically promote immune response against tumor, the minus-strand RNA viral vector expressing one or more tumor antigens is introduced into dendritic cells, or dendritic cells
5 activated by the minus-strand RNA viral vector are pulsed with tumor antigens. The tumor antigens may be tumor cell-specific antigens (*i.e.*, existing in tumor cells but absent in non-tumor cells) or antigens that are expressed at a higher level in tumor cells than in non-tumor cells of the same type. The immune system is stimulated through the administration of the dendritic cells. When CTL acts as a major effector, a desired intercellular or extracellular tumor antigen can be
10 used. When an antibody is reacted as the effector by using dendritic cells to activate CD4 T cells which triggers the induction of antibody production through B cell activation, it is preferred to use an antigen presented on the cell surface. For example, a cell surface receptor or cell adhesion protein can be used as the antigen. The tumor antigens include, for example, Muc-1 or Muc-1-like mucin tandem repeat peptide that relate ovarian cancer or the like (U.S. Patent No.
15 5,744,144); E6 and E7 proteins of human papilloma virus, which cause cervical cancer; melanoma antigens MART-1, MAGE-1, -2, -3, gp100, and tyrosinase; prostate cancer antigen PSA; as well as CEA (Kim, C. *et al.*, Cancer Immunol. Immunother. 47 (1998) 90-96) and Her2neu (HER2p63-71, p780-788; Eur. J. Immunol. 2000; 30: 3338-3346).

[0092]

20 Dendritic cells that are prepared according to the present invention are useful as effective immunotherapy for cancers and infectious diseases. Immunological sensitization by dendritic cells introduced with a gene of a tumor antigen or infectious disease-related antigen or T cells stimulated with such dendritic cells serves as an effective method for inducing antitumor or anti-infectious disease immunity in patients.

25 [0093]

The cells can also be applied to general diseases. To treat diabetes, for example, a peptide of an insulin fragment can be used as an epitope in animal models of type I diabetes (Coon, B. *et al.*, J. Clin. Invest., 1999, 104 (2):189-94).

[0094]

30 In addition, by expressing a cytokine in dendritic cells, the cells stimulate the immune system, thereby enhancing immune responses against cancers or infectious microorganisms. Thus, dendritic cells introduced with a gene encoding a cytokine are also useful in the treatment of cancers and other diseases for which cytokine therapy is expected to be effective. For example, immunostimulatory cytokines include interleukins (for example, IL-1alpha, IL-1beta,
35 IL-2, IL-3, IL-4, IL-6, IL-7, IL-8, IL-9, IL-10, IL-12, IL-15, IL-18, IL-19, IL-20, IL-21, IL-23, and IL-27), interferons (for example, IFN-alpha, IFN-beta, and IFN-gamma), tumor necrosis

factor (TNF), transforming growth factor (TGF)-beta, granulocyte colony stimulating factor (G-CSF), macrophage colony stimulating factor (M-CSF), granulocyte macrophage colony stimulating factor (GM-CSF), insulin-like growth factor (IGF)-I, IGF-2, Flt-3 ligand, Fas ligand, c-kit ligand, and other immunomodulatory proteins (such as chemokines and costimulatory molecules).

[0095]

The amino acid sequences of these cytokines are well known to those skilled in the art. One may refer to: for IL-4, for example, Arai *et al.* (1989), J. Immunol. 142 (1) 274-282; for IL-6, for example, Yasukawa *et al.* (1987), EMBO J., 6 (10): 2939-2945; for IL-12, for example, Wolf *et al.* (1991), J. Immunol. 146 (9): 3074-3081; for IFN-alpha, for example, Gren *et al.* (1984) J. Interferon Res. 4 (4): 609-617, and Weismann *et al.* (1982) Princess Takamatsu Symp. 12: 1-22; for TNF, for example, Pennica *et al.* (1984) Nature 312: 724-729; for G-CSF, for example, Hirano *et al.* (1986) Nature 324: 73-76; and for GM-CSF, for example, Cantrell *et al.* (1985) Proc. Natl. Acad. Sci. (USA) 82 (18): 6250-6254. Vectors can be introduced into dendritic cells by designing them to include natural genes encoding these cytokines or mutant genes that still encode functional cytokines due to the degeneracy of genetic code.

[0096]

Moreover, the genes may be modified to express modified forms of the cytokines. For example, a cytokine that has both precursor and matured forms (for example, those producing active fragments by cleavage of their signal peptides, or by restrictive proteolysis) may be genetically modified to express either the precursor or the matured form. Other modified forms (for example, fusion proteins of an active fragment of a cytokine and a heterologous sequence (for example, heterologous signal peptide)) can also be used.

[0097]

Dendritic cells genetically modified as described above are useful to stimulate T cells of patients themselves *in vivo*. In addition, these genetically-modified dendritic cells are also useful to stimulate T cells *in vitro*. A sensitized T cell may be administered to patients to stimulate the patient's immune system by *ex vivo* immunotherapy.

[0098]

The present invention relates to a method for producing T cells stimulated with dendritic cells, which includes the steps of: (a) contacting a minus-strand RNA viral vector with a dendritic cell or a precursor cell thereof; (b) differentiating the cell into mature dendritic cell; and (c) contacting the mature dendritic cell with a T cell. The antigen to be presented by the dendritic cell may be a protein (or a processed product thereof) expressed from the vector or a protein exogenously pulsed into the dendritic cell. The dendritic cell introduced with the minus-strand RNA viral vector activates T cells and induce CTLs.

[0099]

The present invention also relates to a method for stimulating the immune system using a dendritic cell produced by the method of the present invention. For example, patients suffering from infectious disease, cancer, or the like can be treated to stimulate their immune system. This method includes the step of administering a dendritic cell or a T cell. Specifically, the method includes the step of (a) administering into a patient a therapeutically effective amount of dendritic cells introduced with the minus-strand RNA viral vector; or (b) administering into a patient a therapeutically effective amount of T cells stimulated by dendritic cells introduced with the minus-strand RNA viral vector. The minus-strand RNA viral vector may carry genes encoding one or more antigens or cytokines associated with diseases or may carry no foreign gene. Since the minus-strand RNA viral vector activates dendritic cells by infecting the cells, dendritic cells infected with the vector carrying no foreign gene can also activate patients' immune system. More highly effective dendritic cells can be obtained by pulsing dendritic cells with an antigen peptide to present the desired antigen. Alternatively, when T cells are contacted with dendritic cells *in vitro*, it is preferred to collect T cells from a patient and carry out *ex vivo* administration.

[0100]

The appropriate *in vivo* dose of the vector varies depending on the disease, patient's weight, age, sex, and symptom, purpose of administration, form of administered composition, administration method, transfer gene, and the like, but can be appropriately determined by those skilled in the art. The route of administration can be appropriately selected, and includes, for example, percutaneous, intranasal, transbronchial, intramuscular, intraperitoneal, intravenous, intraarticular, and subcutaneous administration. The administration may be local or systemic. It is preferred to administer the vector at a dose within the range of preferably about 10^5 to about 10^{11} CIU/ml, more preferably about 10^7 to about 10^9 CIU/ml, and most preferably about 1×10^8 to about 5×10^8 CIU/ml, in a pharmaceutically acceptable carrier. The amount per dose for human is preferably 2×10^5 to 2×10^{11} CIU, which is administered once or more within a range where the side effects are clinically acceptable. The same applies to the number of doses per day. Regarding nonhuman animals, for example, a dose converted from the above-described dose based on the body weight ratio between the subject animal and human or the volume ratio (e.g., mean value) of the target site for administration. In addition, when it becomes necessary to suppress the proliferation of the transmissible minus-strand RNA viral vector after administration to subjects or cells due to the completion of treatment, through the administration of an RNA-dependent RNA polymerase inhibitor the proliferation of the viral vector can be specifically suppressed without damaging the host.

[0101]

For *ex vivo* administration, the vector is contacted with dendritic cells *ex vivo* (for example, in a test tube or dish). It is preferable to administer at a MOI of between 1 to 500, more preferably 2 to 300, even more preferably 3 to 200, still more preferably 5 to 100, and yet more preferably 7 to 70. The subject to which the vector is administered is not particularly
5 limited, and includes, for example, birds and mammals, including chicken, quail, mouse, rat, dog, pig, cat, bovine, rabbit, sheep, goat, monkey, and human, and other vertebrates.

[0102]

When administering a dendritic cell introduced with the vector, the cell can be generally injected by intramuscular, intraperitoneal, subcutaneous, or intravenous injection, or direct
10 injection into the lymph node. Preferably, the cell is administered into patients by subcutaneous or intraperitoneal injection, or direct injection into the lymph node. Patients can be administered typically with 10^5 to 10^9 transformed dendritic cells, preferably 10^6 to 10^8 cells, and more preferably about 10^7 cells.

[0103]

15 When T cells activated with the dendritic cells are administered, for example, the T cells can be administered at a dose of about 10^5 to 10^9 cells, preferably 10^6 to 10^9 cells, and more preferably 10^8 to 10^9 cells per 1m^2 body surface area by intravenous injection (see Ridell *et al.*, 1992, Science 257: 238-241). The injection can be repeated at desired intervals (for example, monthly). After the administration, recipients may be monitored for any side effects during or
20 after T cell injection, if required. In this case, it is preferred that T cells are obtained from the same patient from whom the dendritic cells have been derived. Alternatively, the T cells may be collected from a patient, while the dendritic cells to stimulate the T cells may be derived from an HLA-compatible healthy donor. Conversely, the dendritic cells may be collected from a patient, while the T cells may be derived from an HLA-compatible healthy donor.

25 [0104]

Cells containing the dendritic cells as the active ingredient of vaccines that are produced according to the present invention are inoculated as therapeutic vaccines to the human body. Thus, it is safer to make them deficient in growth capacity. For example, it is known that the growth capacity of cord blood-derived monocytes is extremely reduced after the induction of
30 differentiation. However, to use the cells as safer cell vaccines, the growth capacity can be reduced or eliminated without losing the vaccine function by treating the cells with heat, radiation, mitomycin C, or the like. For example, when X-ray irradiation is used, X-ray can be irradiated at a total radiation dose of 1,000 to 3,300 Rad. With regard to the mitomycin C treatment, mitomycin C can be added to the dendritic cells at a concentration of 25 to 50 $\mu\text{g/ml}$
35 and incubated at 37°C for 30 to 60 minutes. When the cells are treated with heat, for example, the cells can be subjected to a heat treatment of 50 to 65°C for 20 minutes.

[Examples]

[0105]

Hereinbelow, the present invention is specifically described in the context of Examples; however, it is not to be construed as being limited thereto. All publications cited herein are
5 incorporated as a part of the specification.

[0106]

A. Examination on introduction efficiency:

[Experiment 1]

Monocytes from healthy donors were enriched through negative selection.
10 RosetteSepTM-human monocyte enrichment cocktail (Stem Cell Technology Inc.) was used in the negative selection to enrich the monocytes. Specifically, a tetrameric antibody (consisting of two antibody molecules linked together; one is anti-glycophorin A antibody that recognizes erythrocytes, and the other an antibody that recognizes a surface antigen of mononuclear cells) was used to bind cells to be removed to erythrocytes, and the cells were removed using Ficoll
15 PaqueTM Plus (Pharmacia Biotech Inc.). Through negative selection, cells expressing CD2, CD3, CD8, CD19, CD56, and CD66b were eliminated, and the remaining cells were used as monocyte-enriched cells in the following induction of DC differentiation. At this stage, 65-80% were CD14⁺ cells. GM-CSF (500 U/ml) and IL-4 (250 U/ml) were added to the monocyte-enriched cells, and the cells were cultured in endotoxin-free RPMI supplemented with
20 10% FCS to prepare DCs. After 3 to 4 days, half of the culture supernatant was exchanged with fresh culture medium having the same composition. The cells were confirmed to be positive in the expression of costimulatory molecules, and CD11c, HLA-class II (DR, DP, and DQ), and CD1a, and not to present other lineage markers (CD3, CD56, CD19, CD15, and CD14) (Fig. 1, and data not shown). These cells were used to test the efficiency of vector introduction.
25 At this stage, 90 to 98% of the viable cells expressed DC markers (CD11c, and HLA-class II (DR, DP, and DQ)).

Although the above-described kit was used in the selection in this Example, a similar selection can also be performed by using antibody-coated magnetic beads. The use of the beads is preferred to preparing cells on a large scale, such as to collect mononuclear cells through
30 blood cell separation or the like.

[0107]

[Experiment 2]

Sendai virus vector (SeV-GFP) (propagative; WO 00/70070) that expresses the green fluorescent protein (GFP) was infected to the DCs obtained in Experiment 1 (7 days after
35 differentiation induction) at various MOIs. Changes in the cell count, the expression level of GFP, and the expression levels of costimulatory molecules were investigated over time. The

result showed that %GFP reached the maximal level when the MOI was 20 or greater (Figs. 2 to 5). The mean fluorescence intensity (MFI) of GFP can be further increased when the MOI is increased up to 100 (data not shown). Further, the MFI of GFP increased up to day 8. The level of costimulatory molecules (CD80 and CD86) as a whole became maximal when the MOI was 20 or greater. Regarding the decrease in cell count, less change was observed within the MOIs of 1 to 20, and a slight decrease was observed at a MOI of 50 without significant difference (Fig. 6).

[0108]

[Experiment 3]

The DCs were infected with SeV-GFP at a MOI of 20, and the expression of GFP was examined over time using FACS. As a result, the expression decreased after two weeks (the cell count was also decreased) but GFP expressing cells were detectable up to two months later (Fig. 7). As described in the Example below, DCs are activated by the infection of the minus-strand RNA viral vector. Thus, gene transfer into DCs using the minus-strand RNA viral vector is clinically applicable to vaccination. The administration can be achieved *in vivo* or *ex vivo*. The gene expression can be maintained in the body for a long period by, for example, frequently administering DCs infected with the vector through *ex vivo* administration.

[0109]

[Experiment 4]

The activation and infection efficiency were examined. It was examined whether the efficiency of vector infection was altered by the activation. DCs cultured for 7 days were stimulated with LPS (1 μ g/ml) for two days, infected with SeV-GFP at a MOI of 30, and after 2 days GFP was analyzed by FACS. Alternatively, 2 days after SeV-GFP infection, LPS stimulation (for two days) was carried out under the same condition. (Figs. 8 and 9)

Results: human DCs were found to be nearly 60% positive in %GFP after activation with LPS. In contrast, in mouse DCs, the positivity rate was very low (data not shown). However, MFI was also very low in human, showing a drastic decrease in the efficiency of gene transfer into DCs after activation. In contrast, the efficiency of gene transfer was not altered by LPS stimulation after vector introduction. These results demonstrate that it is preferable to use immature DCs, *i.e.* non-activated DCs, for gene transfer into DCs using the minus-strand RNA viral vector.

[0110]

[Experiment 5]

The contact time required for infection was examined (Fig. 10). The results demonstrate that gene transfer can be achieved within about 30 minutes or less.

[0111]

[Experiment 6]

A previous report described success in producing gene-transferred DCs through the introduction of genes into CD34 cells and the induction of differentiation into DCs (J. Immunol. Meth. 2002; 153-165). A similar method was conducted for SeV-GFP. CD34 positive stem cells (CD34 > 90%) were separated from human cord blood using CD34 microbeads. After infection at a MOI of 0, 10, or 100, the cells were washed well. The cells were cultured in RPMI + 10% FCS supplemented with SCF (50 ng/ml), GM-CSF (500 U/ml), and TNF-alpha (50 ng/ml) for 3 days, then, passaged in a medium supplemented with SCF (50 ng/ml), GM-CSF (500 U/ml), IL-4 (250 U/ml), and TNF-alpha (50 ng/ml) (half of the medium was exchanged every 3 to 4 days), and GFP expression was examined 13 days after the vector infection. As a result, the gene transfer efficiency reached 65 to 70%, and DCs having better expression efficiency of GFP than those prepared with other vectors were prepared. By analyzing the expression of costimulatory molecules, more activated DCs were recovered from the infected DCs than uninfected DCs. (Figs. 11 and 12).

According to the Examples described above, it was demonstrated that the introduction efficiency of the minus-strand RNA virus is considerably higher than that of lentivirus or retrovirus, and an efficiency comparable to that of adenovirus can be achieved rapidly and very easily. In addition, it was found that the activation markers were not altered by using other vectors; however, DC activation can be induced by the infection of the minus-strand RNA virus.

[0112]

B. Evaluation of DC function after introduction

[Experiment 1]

DCs were infected with SeV-GFP at a MOI of 30 to 50. On the following day, the cells were stimulated by LPS (for 2 days) and tested for the expression of costimulatory molecules.

As controls, the conditions of LPS stimulation alone, SeV-GFP infection alone, and no LPS stimulation nor SeV-GFP infection were examined and compared.

Results: The obtained results demonstrate that DC activation occurs by SeV infection alone.

Comparable to LPS: CD80(+) HLA-DR(-) CD83(-)

Higher than LPS: CD86(+) CCR7(-)

Lower than LPS: CD40(-)

(+) indicates where synergistic effect can be obtained by using LPS and SeV. (Figs. 13 to 15)

[0113]

[Experiment 2]

DCs were infected with SeV-GFP at a MOI of 30 (some groups were stimulated with LPS on the next day of the infection or 3 days after the infection)). The phagocytic activity was

examined in the groups similarly to those as described in Experiment 1 (1 μ m PCV-RED latex-microspheres were used. The bar graphs represent the activity after subtraction of positive background at 4°C).

Results: The phagocytic activity was found to be reduced in the cells infected with SeV due to the activation as was shown by activation markers. In particular, the higher the expression level of GFP, the lower the phagocytic activity. Thus, for example, when a tumor cell lysate is used to present tumor antigens on DCs, it is preferred to co-culture DCs with the lysate before the introduction of the minus-strand RNA viral vector into DCs. (Figs. 16 to 17)

[0114]

[Experiment 3]

T cell activating ability was examined using the same experimental groups by irradiating the DCs at 3,000 rad. (Purified ($CD3^+ > 95\%$) allo or syngenic T cells were co-cultured with DCs at various DC doses for 3 days). Syngenic T cells were used as an indicator of response to SeV-GFP.

Results: Due to the low DC ratio and the number of T cells, the differences were relatively insignificant. Nonetheless, it was found that SeV infection alone had an allo T cell-stimulating effect equivalent to LPS (Fig. 18). DCs can also be used without irradiation.

[0115]

C. *In vivo* induction of cancer antigen-specific CTLs

Using the method described above in subsection A, $CD14^+$ cells were enriched from human peripheral blood (healthy donors with HLA-A 0201), and immature dendritic cells were prepared using x-vivo 15 (Cambrex) + 2% autologous serum as a medium, supplemented with GM-CSF (500 U/ml), and IL-4 (250 U/ml) (half of the medium was exchanged every 3 to 4 days). The prepared immature dendritic cells were divided into the following three groups, and then further cultured for 48 hours in the presence of GM-CSF (500 U/ml) and IL-4 (250 U/ml):

Group 1: no addition;

Group 2: infected with SeV-GFP (MOI 30); and

Group 3: stimulation by cytokine cocktail (50 ng/ml IL-1 β , 500 ng/ml IL-6, 2,500 U/ml INF- α , 100 ng/ml TNF- α , and 20 μ M PGE2).

Next, dendritic cells were recovered and pulsed with MART-1 peptide (EAAGIGILTV (SEQ ID NO: 9); 50 μ g/ml for 3 hours). T cells in peripheral blood from the same healthy donor from whom the dendritic cells had been obtained were enriched through negative selection ($CD3^+ > 97\%$), and were co-cultured with peptide-pulsed dendritic cells of the above three groups for 7 days (X-vivo 15 + 2% autologous serum). (Half of the medium was exchanged 3 to 4 days or when the medium changed yellow. After the T cells and dendritic cells were co-cultured at the first stimulation, 100 U/ml IL-2 was added at the third day.) This treatment

was repeated twice. The cells were recovered from each mixed culture fluid and used as effector cells in CTL assay.

T2 cells (TAP deficient cell line, a T cell-B cell hybridoma, obtained from a donor with HLA-A2⁺) was used as target cells. These cells lack TAP (the transporter to class I), and therefore are incapable of transferring peptides produced through cytoplasmic proteolysis to class I. Thus, when a peptide is exogenously added, the peptide is loaded onto class I resulting in class I expression. The target cells were pulsed with mutant MART-1 peptide (ELAGIGILTV (SEQ ID NO: 10); a peptide with potentiated HLA-A2 binding ability without any alteration in the T cell receptor recognition site as compared to the peptide used in the above-described stimulation) or with influenza peptide (Flu; a peptide as a third party; GILGFVFTL (SEQ ID NO: 11)), and labeled with Cr. The effector T cells of the above three groups were co-cultured with each of the two types of targets at a ratio of 20:1, 10:1, 5:1, or 2.5:1 for four hours to examine the CTL activity.

The combinations used in the experiment are summarized below.

Effector cells	Target cells	Symbols in the figure
Effector T cells of Group 1	Mutant MART1 peptide + T2 cells	Solid line with closed squares
Effector T cells of Group 2	Mutant MART1 peptide + T2 cells	Solid line with closed triangles
Effector T cells of Group 3	Mutant MART1 peptide + T2 cells	Solid line with closed inverted triangles
Effector T cells of Group 1	Flu peptide + T2 cells	Dotted line with closed diamonds
Effector T cells of Group 2	Flu peptide + T2 cells	Dotted line with closed circles
Effector T cells of Group 3	Flu peptide + T2 cells	Dotted line with open squares

Results: MART-1 specific CTL cannot be induced when the T cells are stimulated by the non-activated DCs (MART1 peptide +) among the three groups described above. However, as a positive control, when T cells were stimulated using dendritic cells that were activated by cytokines (a method which most intensively activates cells among the current dendritic cell therapy for antitumor immunity), MART-1 specific CTLs could be induced (a similar result was obtained when, instead of the mutant MART-1 peptide, the MART-1 peptide as used in the stimulation was used to pulse the target). When dendritic cells introduced with genes using SeV were used, a CTL activity comparable to the positive control was obtained. Specifically,

when determined by the CTL assay, it was shown that dendritic cells were activated by SeV infection alone, and that they can induce CTLs *in vitro* to the same level as dendritic cells activated by cytokines. When SeV is used for T cell activation, the activation can be achieved simultaneously with the introduction of the target gene, which makes it unnecessary to add
5 activation factors, such as cytokines, and thus contributes to cost reduction, time saving, and retaining cell viability.

[Industrial Applicability]

[0116]

The present invention facilitates the efficient introduction of genes into dendritic cells.

10 The vector of the present invention is preferably used to induce protective immunity against viruses, bacteria, and the like, and also in anti-cancer immunotherapy and the like. Since dendritic cells have strong ability to induce immunity, DC vaccine that induces antigen-specific cellular immunity can be produced by introducing a desired antigen gene or immune-activating gene into dendritic cells using the method of the present invention.

15 [Brief Description of the Drawings]

[0117]

[Fig. 1] Figure depicts graphs showing phenotypes of dendritic cells derived from mononuclear cells in monocyte-enriched peripheral blood cells. Viable cells recognized by PI were gated, and the expression of CD11c and HLA-class II (DR, DP, and DQ) was observed
20 using anti-CD11c-PE-conjugated antibody and anti-HLA-class II (DR, DP, and DQ) FITC-conjugated antibody (panel A). Furthermore, a gate was selected for cells positive for both CD11c and HLA-class II (DR, DP, and DQ), and the expression levels detected with: (1) anti-CD14-APC-conjugated antibody; (2) anti-CD1a-APC-conjugated antibody; and (3) anti-CD80-biotin-conjugated antibody (secondarily stained with streptavidin-APC) relative to
25 that of CD11c are shown in dot plots (panel B to D). In the Examples, "Class II" indicates a result obtained using an antibody recognizing all of HLA-DR, DQ, and DP, and "HLA-DR" indicates a result obtained using an antibody specifically recognizing HLA-DR.

[Fig. 2] Figure depicts graphs showing the expression of GFP and costimulatory molecules in DCs introduced with SeV-GFP.

30 [Fig. 3] Figure depicts graphs showing the introduction efficiency of SeV-GFP into human monocyte-derived dendritic cells and the activation of the dendritic cells (day 2 after infection).

[Fig. 4] Figure depicts graphs showing the introduction efficiency of SeV-GFP into human monocyte-derived dendritic cells and the activation of the dendritic cells (day 4 after
35 infection).

[Fig. 5] Fig. 5 depicts graphs showing the introduction efficiency of SeV-GFP into

human monocyte-derived dendritic cells and the activation of the dendritic cells (day 8 after infection).

[Fig. 6] Figure depicts a graph showing alterations in DC count after SeV-GFP introduction.

5 [Fig. 7] Figure depicts graphs showing the duration of GFP expression after SeV-GFP introduction.

[Fig. 8] Figure depicts graphs showing the effect of LPS stimulation on the introduction efficiency of SeV-GFP into human DCs.

10 [Fig. 9] Figure depicts graphs showing the effect of LPS stimulation on the introduction efficiency of SeV-GFP into human DCs.

[Fig. 10] Figure depicts graphs showing the result of examination on the incubation time for gene transfer into DCs.

[Fig. 11] Figure depicts graphs showing gene transfer into DCs derived from cord blood.

15 [Fig. 12] Figure depicts graphs showing gene transfer into DCs derived from cord blood.

[Fig. 13] Figure depicts graphs showing the expression of costimulatory molecules after gene transfer (as compared with LPS stimulation).

20 [Fig. 14] Figure depicts graphs showing the expression of costimulatory molecules after gene transfer (as compared with LPS stimulation).

[Fig. 15] Figure depicts graphs showing the expression of costimulatory molecules after gene transfer (as compared with LPS stimulation).

[Fig. 16] Figure depicts graphs showing the phagocytic ability after gene transfer.

[Fig. 17] Figure depicts graphs showing the phagocytic ability after gene transfer.

25 [Fig. 18] Figure depicts graphs showing the allo-T-cell stimulating ability of DCs introduced with SeV-GFP.

[Fig. 19] Figure shows the results of *in vitro* induction of MART-1 specific CTLs by introducing the minus-strand RNA viral vector.

[Sequence Listing]

SEQUENCE LISTING

<110> Dनावेक रीसर्च इं.क.

5

<120> Method for producing transgenic dendritic-cells

<130> D3-A0307

10

<140>

<141>

<160> 11

15

<170> PatentIn version 3.1

<210> 1

<211> 10

<212> DNA

20

<213> Artificial

<220>

<223> artificially synthesized sequence

25

<400> 1

ctttcacct

10

<210> 2

30

<211> 15

<212> DNA

<213> Artificial

<220>

35

<223> artificially synthesized sequence

<400> 2

tttttcttac tacgg

15

5 <210> 3

<211> 18

<212> DNA

<213> Artificial

10 <220>

<223> artificially synthesized sequence

<400> 3

cggccgcaga tcttcacg

18

15

<210> 4

<211> 18

<212> DNA

20 <213> Artificial

<220>

<223> artificially synthesized sequence

25 <400> 4

atgcatgccg gcagatga

18

<210> 5

30 <211> 18

<212> DNA

<213> Artificial

<220>

35 <223> artificially synthesized sequence

<400> 5

gttgagtact gcaagagc

18

5 <210> 6

<211> 42

<212> DNA

<213> Artificial

10 <220>

<223> artificially synthesized sequence

<400> 6

tttgccggca tgcattgttc ccaaggggag agttttgcaa cc

42

15

<210> 7

<211> 18

<212> DNA

20 <213> Artificial

<220>

<223> artificially synthesized sequence

25 <400> 7

atgcatgccg gcagatga

18

<210> 8

30 <211> 21

<212> DNA

<213> Artificial

<220>

35 <223> artificially synthesized sequence

<400> 8

tgggtgaatg agagaatcag c

21

5 <210> 9

<211> 10

<212> PRT

<213> Artificial

10 <220>

<223> an artificially synthesized peptide

<400> 9

15 Glu Ala Ala Gly Ile Gly Ile Leu Thr Val

1

5

10

<210> 10

20 <211> 10

<212> PRT

<213> Artificial

<220>

25 <223> an artificially synthesized peptide

<400> 10

Glu Leu Ala Gly Ile Gly Ile Leu Thr Val

30

1

5

10

<210> 11

<211> 9

35 <212> PRT

<213> Artificial

<220>

<223> an artificially synthesized peptide

5 <400> 11

Gly Ile Leu Gly Phe Val Phe Thr Leu


1

5


10

[Document Name] Drawings


[Fig. 1]




[Fig. 2]




[Fig. 3]




[Fig. 4]




[Fig. 5]




[Fig. 6]




[Fig. 7]




[Fig. 8]




[Fig. 9]




[Fig. 10]







[Fig. 11]




[Fig. 12]




[Fig. 13]




[Fig. 14]




[Fig. 15]




[Fig. 16]




[Fig. 17]



[Fig. 18]



[Fig. 19]



[Document Name] Abstract

[Abstract]

[Problems to Be Solved] The present invention provides a method for producing a gene transferred dendritic cell.

- 5 [Means for Solving the Problems] The present invention provides a method for introducing a gene into a dendritic cell, which includes the step of contacting a minus-strand RNA virus with a dendritic cell. The present invention also provides a method for producing a gene transferred dendritic cell, which includes the step of contacting a minus-strand RNA virus with a dendritic cell. The present invention also provides gene transferred dendritic cells produced by such
- 10 methods. Furthermore, the present invention provides a method for activating a dendritic cell, which includes the step of contacting a minus-strand RNA virus with a dendritic cell. The present invention enables efficient gene delivery into dendritic cells. Dendritic cells introduced with antigen gene or cytokine gene are useful as vaccine.

[Selected Drawings] None

15